

- 1 Express $\frac{12}{x+1} - (7-x)$ as a single simplified fraction. [1]

Without using a calculator, solve $\frac{12}{x+1} \leq 7-x$. [3]


- 2 (i) Find $\frac{d}{dx} \tan^{-1}(x^2)$. [1]

(ii) Hence, or otherwise, evaluate $\int_0^1 x \tan^{-1}(x^2) dx$ exactly. [3]

- 3 (i) Find $\frac{d}{dx}(3x^2 2^x)$. [2]

(ii) Find the equation of the tangent to the curve $y = 3x^2 2^x$ at the point where $x = 1$, giving your answer in exact form. [3]

- 4 The graph for $y = f(x)$ is given below, where $y = 10 - x$, $y = 6$ and $x = 4$ are asymptotes. The turning points are $(-3, 5)$ and $(6, 0)$, and the graph intersects the y -axis at $(0, 6)$.



On separate diagrams, sketch the graphs of

- (i) $y = f(|x|)$, [3]

(ii) $y = \frac{1}{f(x)}$. [3]

5 Referred to the origin O , points P and Q have position vectors $3\mathbf{a}$ and $\mathbf{a} + \mathbf{b}$ respectively. Point M is a point on QP extended such that $PM:QM$ is 2:3.

(i) Find the position vector of point M in terms of \mathbf{a} and \mathbf{b} . [2]

(ii) Find $\overrightarrow{PQ} \times \overrightarrow{OM}$ in terms of \mathbf{a} and \mathbf{b} . [3]

(iii) State the geometrical meaning of $\frac{|\overrightarrow{PQ} \times \overrightarrow{OM}|}{|\overrightarrow{PQ}|}$. [1]

6 A curve C has equation $y = f(x)$, where the function f is defined by

$$f : x \mapsto \frac{12 - 3x}{x^2 + 4x - 5}, \quad x \in \mathbb{R}, x \neq -5, x \neq 1.$$

(i) Find algebraically the range of f . [3]

(ii) Sketch C , indicating all essential features. [4]

(iii) Describe a pair of transformations which transforms the graph of C on to the graph of $y = \frac{9 - x}{x^2 - 6x}$. [2]

7 Given that $\sin^{-1} y = \ln(1 + x)$, where $0 < x < 1$, show that $(1 + x) \frac{dy}{dx} = \sqrt{1 - y^2}$. [2]

(i) By further differentiation, find the Maclaurin expansion of y in ascending powers of x , up to and including the term in x^2 . [4]

(ii) Use your expansion from (i) and integration to find an approximate expression for $\int \frac{\sin(\ln(1 + x))}{x} dx$. Hence find an approximate value for $\int_0^{0.5} \frac{\sin(\ln(1 + x))}{x} dx$. [3]

[Turn over

- 8 (a) A sequence of numbers $a_1, a_2, a_3, \dots, a_{64}$ is such that $a_{n+1} = a_n + d$, where $1 \leq n \leq 63$ and d is a constant. The 64 numbers fill the 64 squares in the 8×8 grid in such a way that a_1 to a_8 fills the first row of boxes from left to right in that order. Similarly, a_9 to a_{16} fills the second row of boxes from left to right in that order.

1 st column	↓								
1 st row	→	a_1	a_2	a_3	a_4	a_5	a_6	a_7	a_8
		a_9	a_{10}	...					a_{16}
		⋮							

Given that the sums of the numbers in the **first row** and in the **third column** are 58 and 376 respectively, find the values of a_1 and d . [4]

- (b) A geometric series has first term a and common ratio r , where a and r are non-zero. The sum to infinity of the series is 2. The sum of the six terms of this series from the 4th term to the 9th term is $-\frac{63}{256}$. Show that $512r^9 - 512r^3 - 63 = 0$.

Find the two possible values of r , justifying the choice of your answers. [5]

- 9 One of the roots of the equation $z^3 - az - 66 = 0$, where a is real, is w .

(i) Given that $w = b - \sqrt{2}i$, where b is real, find the exact values of a and $\frac{w}{w^*}$. [6]

(ii) Given instead that $w = re^{i\theta}$, where $r > 0$, $-\pi < \theta < -\frac{3\pi}{4}$, find $|aw^2 + 66w|$ and $\arg(aw^2 + 66w)$ in terms of r and θ . [4]

- 10 The point M has position vector relative to the origin O , given by $6\mathbf{i} - 5\mathbf{j} + 11\mathbf{k}$. The line l_1 has equation $x - 7 = \frac{y}{3} = \frac{z + 2}{-2}$, and the plane π has equation $4x - 2y - z = 30$.

(i) Show that l_1 lies in π . [2]

(ii) Find a cartesian equation of the plane containing l_1 and M . [3]

The point N is the foot of perpendicular from M to l_1 . The line l_2 is the line passing through M and N .

(iii) Find the position vector of N and the area of triangle OMN . [5]

(iv) Find the acute angle between l_2 and π , giving your answer correct to the nearest 0.1° . [3]

- 11** [It is given that the volume of a cylinder with base radius r and height h is $\pi r^2 h$ and the volume of a cone with the same base radius and height is a third of a cylinder.]

A manufacturer makes double-ended coloured pencils that allow users to have two different colours in one pencil. The manufacturer determines that the shape of each coloured pencil is formed by rotating a trapezium $PQRS$ completely about the x -axis, such that it is a solid made up of a cylinder and two cones. The volume, $V \text{ cm}^3$, of the coloured pencil should be as large as possible.

It is given that the points P , Q , R and S lie on the curve $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$, where a and b are positive constants. The points R and S are $(-a, 0)$ and $(a, 0)$ respectively, and the line PQ is parallel to the x -axis.

- (i) Verify that $P(a \cos \theta, b \sin \theta)$, where $0 < \theta < \frac{\pi}{2}$, lies on the curve $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$. Write down the coordinates of the point Q . [2]
- (ii) Show that V can be expressed as $V = k\pi \sin^2 \theta (2 \cos \theta + 1)$, where k is a constant in terms of a and b . [3]
- (iii) Given that $\theta = \theta_1$ is the value of θ which gives the maximum value of V , show that θ_1 satisfies the equation $3 \cos^2 \theta + \cos \theta - 1 = 0$. Hence, find the value of θ_1 . [4]

At $\theta = \frac{\pi}{6}$, the manufacturer wants to change one end of the coloured pencil to a rounded-end eraser. The eraser is formed by rotating the arc PS completely about the x -axis.

- (iv) Find the volume of the eraser in terms of a and b . [3]

- 12** A ball-bearing is dropped from a point O and falls vertically through the atmosphere. Its speed at O is zero, and t seconds later, its velocity is $v \text{ ms}^{-1}$ and its displacement from O is $x \text{ m}$. The rate of change of v with respect to t is given by $10 - 0.001v^2$.

- (i) Show that $v = 100 \left(\frac{e^{\frac{t}{5}} - 1}{e^{\frac{t}{5}} + 1} \right)$. [4]

- (ii) Find the value of v_0 , where v_0 is the value approached by v for large values of t . [1]

- (iii) By using chain rule, form an equation relating $\frac{dx}{dt}$, $\frac{dv}{dt}$ and $\frac{dv}{dx}$. Given that $v = \frac{dx}{dt}$, form a differential equation relating v and x . Show that

$$v = 100 \sqrt{1 - e^{-\frac{x}{500}}}. \quad [5]$$

- (iv) Find the distance of the ball-bearing from O after 5 seconds, giving your answer correct to 2 decimal places. [3]

[Turn over

Section A: Pure Mathematics [40 Marks]

- 1 Express $2 \cos \theta \sin \frac{\theta}{2}$ in the form $\sin a\theta - \sin b\theta$, where a and b are constants to be found. [2]

Hence, find the exact value of α , where $0 < \alpha < \pi$, for which

$$\int_{\alpha}^{\pi} \left(3 \cos \frac{3\theta}{2} - \cos \frac{\theta}{2} \right) e^{\cos \theta \sin \frac{\theta}{2}} d\theta = 4 \left(\frac{1}{e} - 1 \right). \quad [5]$$

- 2 (i) Show that $\frac{2}{2r+1} - \frac{3}{2r+3} + \frac{1}{2r+5} = \frac{Ar+B}{(2r+1)(2r+3)(2r+5)}$, where A and B are constants to be found. [2]

- (ii) Hence find $\sum_{r=1}^n \frac{2r+9}{(2r+1)(2r+3)(2r+5)}$. (There is no need to express your answer as a single algebraic fraction.) [4]

- (iii) It is given that $\sum_{r=1}^n \frac{2r+9}{(2r+1)(2r+3)(2r+5)}$ is within 0.01 of the sum to infinity.

Write down an inequality in terms of n , and hence find the smallest possible value of n . [3]

- 3 The function f is defined by

$$f : x \mapsto \frac{-x^2 + 5x - 11}{x - 2}, \quad x \in \mathbb{R}, x \neq 2.$$

- (i) Find the equations of the asymptotes of the curve $y = f(x)$. [3]
- (ii) Determine whether f has an inverse, justifying your answer. [2]

Given that the function g is defined by

$$g : x \mapsto f(x), \quad x \in \mathbb{R}, 2 < x \leq 4,$$

find $g^{-1}(x)$ and state the domain of g^{-1} . [4]

Sketch the graph of $y = g^{-1}g(x)$. [2]

- 4 A curve C has parametric equations

$$x = -\sqrt{t^2 + 4}, \quad y = \frac{\ln t}{t}, \quad \text{where } t > 0.$$

- (i) Show that $\frac{dy}{dx} = \frac{(\ln t - 1)\sqrt{t^2 + 4}}{t^3}$. [3]
- (ii) Find the exact coordinates of the turning point on C , and explain why it is a maximum. [4]
- (iii) Sketch C . [3]
- (iv) Show that the area bounded by C and the lines $x = -\sqrt{13}$ and $x = -\sqrt{5}$ is given by

$$\int_1^3 \frac{\ln t}{\sqrt{t^2 + 4}} dt.$$

Find the area, giving your answer to 4 decimal places. [3]

Section B: Probability and Statistics [60 Marks]

- 5 Mr and Mrs Lee participate in a game show, together with 3 other men and 5 other women. In the first round, the 10 participants are grouped into 5 pairs.
- (i) Find the number of ways the pairings can be done if there is only 1 pair of the same gender. [3]
- After the first round, Mr and Mrs Lee are both eliminated. The remaining 8 participants are seated around a round table. Find the number of ways this can be done if
- (ii) there are no restrictions, [1]
- (iii) the 3 men are not all seated together. [3]

[Turn over

- 6 As the use of email becomes more prevalent, the number of unsolicited email (also known as spam) received increases. Besides advertisements, spam can now be cleverly disguised as business emails and contain malware. Hence, there is a need to use spam filter.


The probability that Yip receives a spam email is p . He uses a spam filter, Spam Guard Plus, to filter his emails. He has the following information:

$$P(\text{an email is classified correctly}) = \frac{41}{50};$$

$$P(\text{an email is classified correctly} \mid \text{it is classified as spam}) = \frac{38}{45};$$

$$P(\text{an email is classified correctly} \mid \text{it is a spam email}) = \frac{19}{20}.$$

- (i) Show that $p = \frac{4}{5}$. Hence, complete the probability tree below. [5]



- (ii) Andy and Betty notices that, on average, 30% and 70% of their emails are spam respectively. State whether Spam Guard Plus would be **(a)** more appropriate for Andy, **(b)** more appropriate for Betty, or **(c)** just as appropriate for both Andy and Betty. Justify your answer. [2]

- 7 Seven members of the school cross country team undergo a new training programme to improve their fitness. During a particular session, each of them has to complete a 200 metre run and to achieve as many push-ups as possible in one minute. The times taken for the 200 metre run, t seconds, together with the number of push-ups each runner achieves, n , are shown in the table.

Student	A	B	C	D	E	F	G
t	38.3	42.1	35.1	40.1	32.0	31.6	41.0
n	44	35	48	42	49	49	40

- (i) Draw a scatter diagram to illustrate the data, labelling the axes. [1]
- (ii) Explain using your scatter diagram why the linear model $n = at + b$ would not be appropriate. [1]

It is thought that the relationship between n and t can be modelled by one of the formulae

$$n = c(t - 30)^2 + d \quad \text{or} \quad n = e(t - 30)^3 + f$$

where c , d , e and f are constants.

- (iii) The product moment correlation coefficient between n and $(t - 30)^2$ is -0.980 , correct to 3 decimal places. Determine, with a reason, which of the 2 models is more appropriate. [2]
- (iv) It is known that student H is able to do 48 push-ups in one minute. It is required to estimate student H's timing for the 200 metre run. Find the equation of a suitable regression line, and use it to find the required estimate. Comment on the reliability of this estimate. [5]
- 8 A bag contains two balls numbered 3, n balls numbered 2 and three balls numbered 1. A player picks two balls at random from the bag at the same time.

If the difference between the numbers on the two balls is 2, the player receives \$6.

If the difference between the numbers on the two balls is 1, the player does not receive or lose any money.

If the numbers on the 2 balls are the same, the player loses \$1.

- (i) Show that the largest value of n such that player is expected to receive money from this game is 8. [5]

For the rest of this question, take the value of n to be 8.

- (ii) Show that the probability that a player loses money in a game is $\frac{16}{39}$. [1]

Victoria plays this game 50 times.

- (iii) Find the probability that she lost money for at least 20 games. [2]
- (iv) The probability that Victoria loses money in r games is more than 0.1. Find the set of values of r . [3]

[Turn over

- 9 Exposure to Volatile Organic Compounds (VOCs), which have been identified in indoor air, is suspected as a cause for headaches and respiratory symptoms. Indoor plants have not only a positive psychological effect on humans, but may also improve the air quality. Certain species of indoor plants were found to be effective removers of VOCs.

A commonly known VOC is Benzene. The following data gives the benzene levels, x (in ppm) in 40 test chambers containing the indoor plant *Epipremnum aureum*.

$$n = 40, \quad \sum (x - 26.0) = -30.1, \quad \sum (x - 26.0)^2 = 214.61.$$

The initial mean Benzene level (in ppm) without *Epipremnum aureum* was found to be 26.0.

- (i) Test, at the 5% level of significance, the claim that the mean Benzene level, μ (in ppm), has decreased as a result of the indoor plant *Epipremnum aureum*. You should state your hypotheses clearly. [5]
- (ii) State, giving a reason, whether there is a need to make any assumptions about the population distribution of the Benzene level in order for the test to be valid. [2]

The Benzene levels of another 50 test chambers containing the indoor plant *Epipremnum aureum* were recorded, The sample mean is \bar{x} ppm and the sample variance is 8.33 ppm².

- (iii) The acceptance region of a test of the null hypothesis $\mu = 26.0$ is $\bar{x} > 25.1$. State the alternative hypothesis and find the level of significance of the test. [4]
- (iv) If the null hypothesis is $\mu = \mu_0$, where $\mu_0 > 26.0$, would the significance level of a test with the same acceptance region in part (iii) be larger or smaller than that found in part (iii)? Give a reason for your answer. [2]

- 10 In this question, you should state clearly the values of the parameters of any distribution you use.**

A bus service plies from a point A in the city, through a point B , and then to its terminal station at point C .

Journey times in minutes from A to B have the distribution $N(28, 4^2)$.

- (i) Find the probability that a randomly selected bus journey from A to B is completed within 35 minutes. [1]

The journey times in minutes from A to C , have the distribution $N(46.2, 4.8^2)$.

- (ii) The journey times in minutes from B to C have the distribution $N(\mu, \sigma^2)$. Given that the journey times from A to B are independent of the journey times from B to C , find the value of μ and show that $\sigma^2 = 7.04$. [3]

- (iii) Find the set of values of k such that at least 90% of all journey times from A to C can be completed within k minutes. [2]

The performance of the bus operation is deemed as “unreliable” if a random sample of 70 journeys from A to C yields a mean journey time exceeding 47 minutes.

- (iv) Two independent random samples of 70 journeys from A to C are taken. Find the probability that both samples will result in the performance of the bus operation to be deemed as “unreliable”. [3]

To improve the reliability performance of the bus operation, more bus lanes are introduced and some bus stops along the bus route are removed. The journey times from A to B are now reduced by 10%, and the journey times from B to C now have the distribution $N(\mu - 1, 8)$.

- (v) Find the probability that two journeys from A to C are completed within a total of 90 minutes. [4]

[Turn over