

Name		( )	Class	
------	--	-----	-------	--



**南 华 中 学**  
**NAN HUA HIGH SCHOOL**  
**PRELIMINARY EXAMINATION 2019**

**Subject : Additional Mathematics**  
**Paper : 4047/01**  
**Level : Secondary Four Express**  
**Date : 29 August 2019**  
**Duration : 2 hours**

**READ THESE INSTRUCTIONS FIRST**

Write your name, class and index number on all the work you hand in.

Write in dark blue or black pen.

You may use an HB pencil for any diagrams or graphs.

Do not use staples, paper clips, glue or correcting fluid / tape.

Answer **all** the questions.

Give non-exact numerical answers correct to 3 significant figures, or 1 decimal place in the case of angles in degrees, unless a different level of accuracy is specified in the question.

The use of an approved scientific calculator is expected, where appropriate.

You are reminded of the need for clear presentation in your answers.

At the end of the examination, fasten all your work securely together.

The number of marks is given in brackets [ ] at the end of each question or part question.

The total number of marks for this paper is 80.

<b>For Examiner's Use</b>

## Mathematical Formulae

### 1. ALGEBRA

#### Quadratic Equation

For the equation  $ax^2 + bx + c = 0$ ,

$$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$

#### Binomial Expansion

$$(a+b)^n = a^n + \binom{n}{1}a^{n-1}b + \binom{n}{2}a^{n-2}b^2 + \dots + \binom{n}{r}a^{n-r}b^r + \dots + b^n$$

where  $n$  is a positive integer and  $\binom{n}{r} = \frac{n!}{r!(n-r)!} = \frac{n(n-1)\dots(n-r+1)}{r!}$

### 2. TRIGONOMETRY

#### Identities

$$\sin^2 A + \cos^2 A = 1$$

$$\sec^2 A = 1 + \tan^2 A$$

$$\operatorname{cosec}^2 A = 1 + \cot^2 A$$

$$\sin(A \pm B) = \sin A \cos B \pm \cos A \sin B$$

$$\cos(A \pm B) = \cos A \cos B \mp \sin A \sin B$$

$$\tan(A \pm B) = \frac{\tan A \pm \tan B}{1 \mp \tan A \tan B}$$

$$\sin 2A = 2 \sin A \cos A$$

$$\cos 2A = \cos^2 A - \sin^2 A = 2 \cos^2 A - 1 = 1 - 2 \sin^2 A$$

$$\tan 2A = \frac{2 \tan A}{1 - \tan^2 A}$$


#### Formulae for $\triangle ABC$

$$\frac{a}{\sin A} = \frac{b}{\sin B} = \frac{c}{\sin C}$$

$$a^2 = b^2 + c^2 - 2bc \cos A$$

$$\Delta = \frac{1}{2} ab \sin C$$

1



The diagram shows a triangle  $ABC$  in which  $AB = 4(\sqrt{3}-1) \text{ cm}$ ,  $BC = 3\sqrt{2}+4 \text{ cm}$ ,  $\angle BAC = 75^\circ$  and  $\angle BCA = \theta$ . Given that  $\sin 75^\circ = \frac{\sqrt{2}}{4}(\sqrt{3}+1)$ , find, without using a calculator, the value of  $\sin \theta$  in the form of  $a + b\sqrt{2}$  where  $a$  and  $b$  are integers. [4]

- 2 Evaluate, without using a calculator,  $\tan\left[\cos^{-1}\left(-\frac{8}{17}\right)\right]$ . [3]

- 3 A function is defined by the equation  $y = \frac{\sin x}{1 - \cos x}$  for  $0 \leq x \leq 2\pi$  where  $x \neq a$ .


- (i) State the exact values of  $a$ . [1]

- (ii) Explain, with reasons, whether the function is increasing or decreasing. [5]

4 A curve has equation  $y = e^{2x-1}$ .

(i) Sketch the graph of  $y = e^{2x-1}$ .

[2]



(ii) The curve  $y = e^{2x-1}$  and  $y = e^{k-x}$  meet at point  $R$  where  $x = 1$ . Find the value of  $k$ . [2]


- 5 (a) Find the range of values of  $n$  for which  $9x^2 + 8nx + 2n^2 > 8$  for all real values of  $x$ .  
[3]

- (b) A curve has equation  $y = (x+3)(x^2 - 3x + 6)$ . Explain why  $y = (x+3)(x^2 - 3x + 6)$  is always positive for  $x > -3$ . [3]



- 6 The coefficient of  $x^3$  in the cubic polynomial  $g(x)$  is  $a$ , where  $a > 0$ . The repeated roots of the equation  $g(x) = 0$  are 2. Find the value of  $a$  if  $g(x)$  has a remainder of  $-\frac{9}{2}$  and 28 when divided by  $(x+1)$  and  $(x-4)$  respectively. [4]

7



The diagram shows a circle with centre  $O$ , diameter  $KL$ .  $NML$  is a tangent to the circle at  $L$  and  $M$  is the midpoint of  $NL$ . The lines  $KN$  and  $OM$  cut the circle at  $P$  and  $Q$  respectively. The lines  $PL$  and  $OQ$  intersect at  $R$ . The line  $LQ$  bisects  $\angle RLM$  and  $\angle NPQ = \angle NKL$ .


- (i) Prove that  $OKPQ$  is a rhombus.

[3]

(ii) Prove that  $KQ \times RQ = LQ \times LR$ .

[3]

8



In the diagram, a surveillance camera is mounted at a point  $A$  that is 20 m above a point  $B$ . A runner runs from point  $B$  along a straight course at a speed of 4 m/s. The surveillance camera tracks the motion of the runner by panning upwards and downwards at point  $A$ . Find the rate of change of the angle that the surveillance camera makes with  $AB$  when the runner is 15 m from  $B$ . Give your answer in radians per second. [5]


- 9 The equation  $2x^2 + x - 4 = 0$  has roots  $\alpha$  and  $\beta$ . The equation  $16x^2 + 21x + p = 0$  has roots  $\frac{1+q\beta^2}{\alpha}$  and  $\frac{1+q\alpha^2}{\beta}$ . Without finding the values of  $\alpha$  and  $\beta$ , find the values of  $p$  and  $q$ .

[6]

10 (i) Find  $\frac{d}{dx}(x^3 \ln x)$ . [2]

(ii) Hence find  $\int x^2 \ln x \, dx$ . [2]

(iii)



The diagram shows the lines  $x = 0.5$ ,  $x = 2$  and part of the curve  $y = x^2 \ln x$ . The curve intersects the x-axis at the point P and the tangent to the curve at P meets the line  $x = 2$  at point Q. Find the total area of the shaded region. [6]




- 11 The table shows experimental values of two variables  $x$  and  $y$ .

$x$	0.1	0.5	1	1.5	2
$y$	-5.95	1.63	0.83	0.61	0.5

It is known that  $x$  and  $y$  are related by the equation  $\frac{\sqrt{x}}{y} = ax + \frac{b\sqrt{x}}{a}$ , where  $a$  and  $b$  are constants.

- (i) Plot  $\frac{1}{y}$  against  $\sqrt{x}$  to obtain a straight line graph. [2]

11(i)




- (ii) Use your graph to estimate the values of  $a$  and  $b$ .

[4]

- (iii) If, instead, a straight line is obtained by plotting  $\frac{1}{y\sqrt{x}}$  against  $\frac{1}{\sqrt{x}}$ , find the gradient of the line.

[2]

12



The diagram shows a kite with vertices  $A(-10, -3)$ ,  $B(0, q)$ ,  $C(2, 1)$  and  $D$ . It is given that angle  $ABO$  is equal to angle  $OBC$ .

- (i) Show that  $q = 2$ .

[4]


- (ii) Find the coordinates of the point  $D$ .

[4]

- (iii) Find the area of the kite  $ABCD$ .

[2]

13




To study the effects of water waves, a wave generator and a rubber duck were placed in a water tank as shown in the diagram above. The height,  $h$  metres, from the bottom of water tank was modelled by  $h = a \sin(kt) + b$ , where  $t$  is the time in hours after midnight and  $a$ ,  $b$  and  $k$  are constants. The motion of the rubber duck was observed for 36 hours. The minimum height of 1.5 m from bottom of water tank was first recorded at 06 00. The maximum height of 2.5 m was first recorded at 18 00.

- (i) Find the values of  $a$ ,  $b$  and  $k$ .

[3]

- (ii) Using the values found in (i), sketch the graph of  $h = a \sin(kt) + b$  for  $0 \leq t \leq 36$ . [2]



- (iii) Find the range of values of  $t$  such that the rubber duck is above 2.1 m. [3]

~ End of Paper ~

Name	( )	Class	
------	-----	-------	--



南 华 中 学

**NAN HUA HIGH SCHOOL**

**PRELIMINARY EXAMINATION 2019**

**Subject : Additional Mathematics**  
**Paper : 4047/02**  
**Level : Secondary Four Express**  
**Date : 2 September 2019**  
**Duration : 2 hours 30 minutes**

**READ THESE INSTRUCTIONS FIRST**

Write your name, class and index number on all the work you hand in.

Write in dark blue or black pen on both sides of the paper.

You may use a soft pencil for any diagrams or graphs.

Do not use staples, paper clips, glue or correcting fluid / tape.

Answer **all** the questions.

Write your answers on the space provided.

Give non-exact numerical answers correct to 3 significant figures, or 1 decimal place in the case of angles in degrees, unless a different level of accuracy is specified in the question.

The use of an approved scientific calculator is expected, where appropriate.

You are reminded of the need for clear presentation in your answers.

The number of marks is given in brackets [ ] at the end of each question or part question.

The total number of marks for this paper is 100.

<b>For Examiner's Use</b>

This paper consists of 24 printed pages.



## Mathematical Formulae

### 1. ALGEBRA

#### Quadratic Equation

For the equation  $ax^2 + bx + c = 0$ ,

$$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$

#### Binomial Theorem

$$(a + b)^n = a^n + \binom{n}{1}a^{n-1}b + \binom{n}{2}a^{n-2}b^2 + \dots + \binom{n}{r}a^{n-r}b^r + \dots + b^n,$$

where  $n$  is a positive integer and  $\binom{n}{r} = \frac{n!}{r!(n-r)!} = \frac{n(n-1)\dots(n-r+1)}{r!}$

### 2. TRIGONOMETRY

#### Identities

$$\sin^2 A + \cos^2 A = 1$$

$$\sec^2 A = 1 + \tan^2 A$$

$$\operatorname{cosec}^2 A = 1 + \cot^2 A$$

$$\sin(A \pm B) = \sin A \cos B \pm \cos A \sin B$$

$$\cos(A \pm B) = \cos A \cos B \mp \sin A \sin B$$

$$\tan(A \pm B) = \frac{\tan A \pm \tan B}{1 \mp \tan A \tan B}$$

$$\sin 2A = 2 \sin A \cos A$$

$$\cos 2A = \cos^2 A - \sin^2 A = 2 \cos^2 A - 1 = 1 - 2 \sin^2 A$$

$$\tan 2A = \frac{2 \tan A}{1 - \tan^2 A}$$

#### Formulae for $\triangle ABC$


$$\frac{a}{\sin A} = \frac{b}{\sin B} = \frac{c}{\sin C}$$

$$a^2 = b^2 + c^2 - 2bc \cos A$$

$$\text{Area of } \triangle ABC = \frac{1}{2} bc \sin A$$

- 1 (a) Without using a calculator, find the value of  $9^x$ , given that  $\frac{1}{3}(5^x)(3^{2x} + 81) = 45^x$ . [4]

- (b) Sketch the graph of  $y = 2\log_4 x - 1$  for  $x > 0$ . [2]



(c) Solve the equation  $\log_3 \frac{1}{9} \sqrt{x} = 1 + 2 \log_x 81$  .

[5]

- 2 In a natural habitat, the population of a certain species of snails is given by  $P = 0.8(Ae^{kt} + 500)$ , where  $A$  and  $k$  are constants and  $t$  is the time in years starting from 1 January 2010. Over a period of 8 years from 1 January 2010 to 31 December 2017, the population decreased from 50 000 to 19 000.

(i) Calculate the values of  $A$  and of  $k$ .

[3]

- (ii) Calculate the year in which the population is 30% more compared to 31 Dec 2017. [3]

- (iii) Explain, with justification, the expected population of the snails over a long period of time. [2]

- 3 (i) Express  $\frac{4x^4 - 4x^3 + 23x^2 - 24x + 5}{x(2x-1)^2}$  in partial fractions.

[5]

(ii) Hence find  $\int \frac{4x^4 - 4x^3 + 23x^2 - 24x + 5}{5x(2x-1)^2} dx$ .

[4]

4 (a) (i) Write down the general term in the binomial expansion of  $\left(\frac{6}{x^2} - \frac{x}{2}\right)^{15}$ . [1]

(ii) Write down the power of  $x$  in this general term. [1]

(iii) Hence, determine the coefficient of  $x^{-9}$  in the expansion of  $\left(\frac{6}{x^2} - \frac{x}{2}\right)^{15}$ . [2]

(iv) Hence determine the coefficient of  $x^{-9}$  in  $\left(\frac{2}{x^2} - \frac{x}{6}\right)^{15}$ . [2]



- (b) The coefficient of  $x^2$  in the expansion, in ascending powers of  $x$ , of  $(1+x)^n (5-2x)^3$  [5]  
is 3210. Find the value of  $n$ , where  $n$  is a positive integer.

- 5 (a) Solve the equation  $4 - |2x + 3| = x$ .

[3]


- (b) A curve has the equation  $y = (2x - 1)^2 - 9$ .

- (i) Explain why the lowest point on the curve has coordinates  $\left(\frac{1}{2}, -9\right)$ .

[1]

- (ii) Sketch the graph of  $y = |(2x-1)^2 - 9|$ .

[3]



- (iii) Determine the set of values of  $m$  such that  $|(2x-1)^2 - 9| = mx - 2$  has no solution.

[2]

6 The coordinates of the points  $A$ ,  $B$  and  $C$  are  $(0, 7)$ ,  $(-1, 0)$  and  $(6, -1)$  respectively.

(i) Show that  $AB$  is perpendicular to  $BC$ .

[2]

(ii) Explain why  $A$ ,  $B$  and  $C$  lie on the circumference of a circle,  $C_1$  with diameter  $AC$ .  
[1]

(iii) Find the centre of  $C_1$ .

[1]

- (iv) The tangent to  $C_1$  at point  $B$  is also a tangent to another circle,  $C_2$ . Given that the centre of  $C_2$ , lies on both the  $y$ -axis and the perpendicular bisector of  $BC$ , find the equation of  $C_2$ . [8]

- 7 (i) Prove that  $\sec 3x(\sin 3x - 2 \sin^3 3x) = \tan 3x \cos 6x$ .


[4]

- (ii) Hence find, for  $0 \leq x \leq \frac{\pi}{3}$ , the values of  $x$  in radians for which

$$-2 \sin \frac{3}{2}x \cos \frac{3}{2}x = \sec 3x (\sin 3x - 2 \sin^3 3x).$$

[6]

8



The diagram shows Car B, which is 48 km due east of Car A. Both cars start moving at the same time. Car A travels due north at a constant speed of 110 km/h while Car B travels due west at a constant speed of 90 km/h.

- (i) The distance between Car A and Car B at time  $t$  hours after the cars started moving is denoted by  $L$  km. Express  $L$  in the form of  $\sqrt{pt^2 + (q - rt)^2}$  where  $p$ ,  $q$  and  $r$  are constants. [3]




- (ii) Given that  $t$  can vary, find the stationary value of  $L$ .

[5]

- (iii) Determine whether this value stationary value of  $L$  gives the maximum or minimum distance between Car  $A$  and Car  $B$ .

[1]

9



For a theatre production, a panel is constructed by joining an isosceles trapezium and a right-angled triangle together.

It is given that  $OA = 5$  m,  $OE = 12$  m and  $OA$  is perpendicular to the base  $OE$ .  $OC$  is perpendicular to  $DE$  and makes an angle  $\theta$  with the base  $OE$ .  $AB$  and  $OC$  are the parallel sides of the trapezium  $OABC$  and  $AB = 2$  m.

The total length of the edges of the panel  $OABCDE$  is represented by  $S$ .

- (i) Show that  $S = 12 \cos \theta + 2 \sin \theta + 22$ .

[3]

- (ii) Express  $S$  in the form of  $R \cos(\theta - \alpha) + Q$ , where  $Q$  is a constant,  $R > 0$  and  $0^\circ < \alpha < 90^\circ$ . [4]

- (iii) A 30 m LED strip is placed along all the edges of the panel  $OABCDE$ . Triangle  $ODE$  of the panel is to be painted in black. Calculate the area to paint. [4]

10

A particle  $X$ , moves in a straight line with velocity,  $v$  m/s, given by  $v = 2t^2 + kt + 63$ , where  $k$  is a constant and  $t$  is the time in seconds, measured from the start of the motion. Its initial displacement from a fixed point  $O$  is  $-8$  m. The minimum velocity of  $X$  occurs at  $t = 5.75$ .

- (i) Find the minimum velocity of  $X$ .

[2]

- (ii) Find the values of  $t$  for which the particle is at instantaneous rest.

[2]

- (iii) Find the distance travelled by particle  $X$  when  $t = 7$ .

[3]

Another particle  $Y$  starts its motion at the same time as particle  $X$  and moves in a straight line with an initial velocity of 6 m/s from  $O$ . Its acceleration,  $a$  m/s<sup>2</sup>, is given by  $a = \frac{3}{5}t$ .

- (iv) Show that particle  $Y$  will not change its direction of motion.

[3]

~End of Paper~



Answer Key


1  $6 - 4\sqrt{2}$

2  $-\frac{15}{8}$

3(i)  $a = 0, 2\pi$

3(ii) Decreasing function

4(i)



4(ii)  $k = 2$

5(a)  $n < -6$  or  $n > 6$

6  $a = \frac{3}{2}$

8  $0.128 \text{ rad / s}$

9  $p = -6, q = -1$


10(i)  $x^2 + 3x^2 \ln x$

10(ii)  $\int x^2 \ln x \, dx = \frac{1}{3}x^3 \ln x - \frac{1}{9}x^3 + c_2$ , where  $c_2$  is an arbitrary constant

10(iii)  $0.639 \text{ unit}^2$  (3 s.f.)



11(i)



11(ii)  $a = 2$  [1.5 to 2.5],  $b = -1.6$  [-2.125 to -1.125]


11(iii)  $-0.8$

12(ii)  $D(1, -1)$

12(iii)  $20 \text{ units}^2$


13(i)  $a = -0.5$ ,  $b = 2$ ,  $k = \frac{\pi}{12}$


13(ii)



13(iii)  $12.8 < t < 23.2$

Answer Key:

1a	40.5
b	
c	6561 or $\frac{1}{9}$
2(i)	-0.123
(ii)	$t = 5.81967$ (must have). In year 2015
(iii)	As $t$ becomes very large (over a long period of time), $e^{-0.122604t}$ approaches 0. Then expected population is $P = 0.8(0 + 500) = 400$
3(i)	$x + \frac{5}{x} + \frac{1}{2x-1} - \frac{3}{(2x-1)^2}$
(ii)	$\frac{1}{10}x^2 + \ln x + \frac{1}{10}\ln(2x-1) + \frac{3}{10(2x-1)} + c$
4a(i)	$\binom{15}{r} (6)^{15-r} \left(-\frac{1}{2}\right)^r (x^{-30+3r})$
a(ii)	Power of $x = -30 + 3r$
a(iii)	-84440070
a(iv)	$-\frac{1430}{243}$
b	9
5a	$x = \frac{1}{3}$ or -7
b(i)	<p>For <math>x \in \mathbb{R}</math>,</p> $(2x-1)^2 \geq 0$ $(2x-1)^2 - 9 \geq -9$ $y \geq -9$ <p>At <math>y = -9, x = \frac{1}{2}</math>. Hence lowest point is <math>\left(\frac{1}{2}, -9\right)</math></p>

b(ii)	
b(iii)	$-2 < m < 1$
6(i)	Product of the gradients of $AB$ and $BC = (7) \times (-\frac{1}{7}) = -1$ $\therefore AB$ is perpendicular to $BC$ .
(ii)	From (i), $AB$ is perpendicular to $BC$ implies that $\angle ABC = 90^\circ$ . Due to $\angle$ in a semi-circle, $AC$ is the diameter of the circle, and $A, B$ and $C$ are points on the circumference of the same circle.
(iii)	(3,3)
(iv)	$x^2 + (y+18)^2 = 100$ .
7(ii)	$0, \frac{\pi}{3}, \frac{\pi}{9}$
8(i)	$L = \sqrt{12100t^2 + (48 - 90t)^2}$
(ii)	37.1
(iii)	$L$ is minimum
9(i)	$S = 12 \cos \theta + 2 \sin \theta + 22$
(ii)	$S = 12.2 \cos(\theta - 9.5^\circ) + 22$
(iii)	$32.2 \text{ m}^2$
10(i)	$-3.125 \text{ m/s}$
(ii)	7 or 4.5
(iii)	117 m
(iv)	For all values of $t$ , since $\frac{3}{10}t^2 > 0$ , $v > 0$ . Particle will not change its direction of motion since velocity is always positive.

