

ZHONGHUA SECONDARY SCHOOL PRELIMINARY EXAMINATION 2020 SECONDARY 4E/5N

Candidate's Name	Class	Register Number

ADDITIONAL MATHEMATICS

4047/01

2 hours

15 September 2020

PAPER 1

Candidates answer on the Question Paper

READ THESE INSTRUCTIONS FIRST

Write your name, class and register number on all the work you hand in. Write in dark blue or black pen on both sides of the paper. You may use an HB pencil for any diagrams or graphs. Do not use staples, paper clips, glue or correction fluid.

Answer **all** the questions.

Give non-exact numerical answers correct to 3 significant figures, or 1 decimal place in the case of angles in degrees, unless a different level of accuracy is specified in the question.

The use of a scientific calculator is expected, where appropriate.

You are reminded of the need for clear presentation in your answers.

The number of marks is given in brackets [] at the end of each question or part question.

The total number of marks for this paper is 80.



Mathematical Formulae

1. ALGEBRA

Quadratic Equation

For the equation $ax^2 + bx + c = 0$,

$$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$

Binomial Theorem

$$(a+b)^{n} = a^{n} + \binom{n}{1}a^{n-1}b + \binom{n}{2}a^{n-2}b^{2} + \dots + \binom{n}{r}a^{n-r}b^{r} + \dots + b^{n},$$

where *n* is a positive integer and
$$\binom{n}{r} = \frac{n!}{(n-r)! r!} = \frac{n(n-1)\dots(n-r+1)}{r!}$$

2. TRIGONOMETRY

Identities

$$\sin^2 A + \cos^2 A = 1.$$
$$\sec^2 A = 1 + \tan^2 A.$$
$$\csc^2 A = 1 + \cot^2 A.$$
$$\sin(A \pm B) = \sin A \cos B \pm \cos A \sin B$$
$$\cos(A \pm B) = \cos A \cos B \mp \sin A \sin B$$
$$\tan(A \pm B) = \frac{\tan A \pm \tan B}{1 \mp \tan A \tan B}$$
$$\sin 2A = 2 \sin A \cos A$$
$$\cos 2A = \cos^2 A - \sin^2 A = 2\cos^2 A - 1 = 1 - 2\sin^2 A$$
$$\tan 2A = \frac{2 \tan A}{1 - \tan^2 A}$$

Formulae for $\triangle ABC$

$$\frac{a}{\sin A} = \frac{b}{\sin B} = \frac{c}{\sin C}$$
$$a^2 = b^2 + c^2 - 2bc \cos A$$
$$\Delta = \frac{1}{2} ab \sin C$$

1 (a) Find the values of x and y which satisfy the equations.

$$3^{x} \times \sqrt{3^{y}} = 1$$

$$4^{x-4} \div 32^{y} = 16^{\frac{1}{x}}$$
 [5]

(b) Without using a calculator, find the values of *a* and *b* such that $7-3\sqrt{3} = (a+b\sqrt{3})(2+\sqrt{3})$, where *a* and *b* are integers.

[4]

2 Find the set of values of the constant k for which the curve $y = (k+2)x^2 - 10x + 2k + 1$ lies completely below the line y = 2x + 3. [4]

- 3 Given that $\sin A = \frac{2}{3}$ and $\tan B = -\frac{1}{\sqrt{3}}$ where angles A and B lie in the same quadrant, leaving your answers in exact form, calculate the value of
 - (a) $\cos 2B$ [2]

(b) $\sec(A-B)$

[3]

- 4 In view of a contagious virus, the government of a particular country has imposed a 'Stay-Home-Notice' on her people to reduce the number of human-to-human transmission cases. It is estimated that the percentage of the population, *P*, complying to the Stay-Home-Notice is given by the equation $P = 100(1 e^{-0.15t})$, where *t* is the number of days after the imposition.
 - (i) Find the percentage of population complying to the 'Stay-Home-Notice' after 5 days of the imposition. [1]

(ii) Find the number of complete days after the imposition that it will take for at least 90% of the population to comply. [2]

(iii) Is it possible for the percentage of this country's population complying to the 'Stay-Home-Notice' to reach 100%? Explain your answer. [1]

5 (i) In the expansion of $\left(2-\frac{x}{3}\right)^n$, show that the ratio of coefficient of the 2nd term to that of the 4th term can be simplified to the expression $\frac{216}{(n-1)(n-2)}$. [4]

(ii) Find the value of n if the ratio in (i) is 108:55.

(iii) Hence, find the term in x^5 .

[2]

[2]

6 The equation of a curve is
$$y = \frac{2x-9}{\sqrt{x^2+1}}$$
.

(i) Show that
$$\frac{dy}{dx} = \frac{9x+2}{\sqrt{(x^2+1)^3}}$$
. [3]

(ii) Find the coordinates of the stationary point of the curve, leaving your answer in exact form. [3]

(iii) Find the nature of this stationary point.

[2]



The diagram shows a triangle *ABC* in which *A* lies on the *y*-axis. The equation of *AB* and *BC* are y = 2x + h and x + 2y = 12h. *M* is midpoint of *AB* and *MC* is parallel to the *x*-axis.

(i) Explain why *AB* is perpendicular to *BC*.

[2]

(ii) Show that coordinates of B is (2h, 5h).

[2]

7

(iii) Express the coordinates of M and of C in terms of h. [3]

(iv) Find the value of h if the area of triangle AMC is 125 square units. [2]

Find the

8

(i) gradient of the tangent at the point where the curve passes through the *x*-axis, [4]

(ii) equation of another tangent to the curve that is parallel to the tangent in (i). [4]

9 The equation of a curve is $y = \cos^3 x + \sin 3x$.

(i) Find expressions for
$$\frac{dy}{dx}$$
 and $\frac{d^2y}{dx^2}$. [4]

(ii) Given that $\frac{d^2 y}{dx^2} + 9y = A\cos x$, find the value of A. [2]

10 The diagram shows an isosceles triangle *RST* with height 12 cm and ST = 10 cm. *PQ* moves towards *ST* at a steady rate of 0.5 cm/s, keeping parallel to *ST*.



If PQ is x cm from R,

(i) Show that
$$PQ = \frac{5x}{6}$$
 cm.

[1]

(ii) Find the area, $A \text{ cm}^2$, of the shaded region in terms of x. [2]

19

[3]

11 It is given that $f(x) = 3\cos 2x - 1$ and $g(x) = \frac{2x}{\pi} - 2$.

- (i) State the least and greatest values of f(x). [2]
- (ii) State the period of f(x). [1]
- (iii) Sketch, on the same axes, the graphs of y = f(x) and y = g(x) for $-\pi \le x \le \pi$. [4]

(iv) State, with detailed workings, the number of solutions of the equation $3\pi \cos 2x = 2x - \pi$ for $-\pi \le x \le \pi$. [2]

12 The function f is defined by $y = (x+1)^3 e^{2x-3}$, for $x > -\frac{5}{2}$, and $x \neq -1$. Explain, with working, whether f is an increasing or decreasing function. [4]



ZHONGHUA SECONDARY SCHOOL PRELIMINARY EXAMINATION 2020

Class

SECONDARY 4E/5N

Candidate's Name

Register Number

ADDITIONAL MATHEMATICS

PAPER 2

4047/02

17 September 2020 2 hours 30 minutes

Candidates answer on the Question Paper

READ THESE INSTRUCTIONS FIRST

Write your name, class and register number on all the work you hand in. Write in dark blue or black pen on both sides of the paper. You may use an HB pencil for any diagrams or graphs. Do not use staples, paper clips, glue or correction fluid.

Answer **all** the questions.

Give non-exact numerical answers correct to 3 significant figures, or 1 decimal place in the case of angles in degrees, unless a different level of accuracy is specified in the question.

The use of a scientific calculator is expected, where appropriate.

You are reminded of the need for clear presentation in your answers.

The number of marks is given in brackets [] at the end of each question or part question.

The total number of marks for this paper is **100**.



Mathematical Formulae

1. ALGEBRA

Quadratic Equation

For the equation $ax^2 + bx + c = 0$,

$$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$

Binomial Theorem

$$(a+b)^{n} = a^{n} + \binom{n}{1}a^{n-1}b + \binom{n}{2}a^{n-2}b^{2} + \dots + \binom{n}{r}a^{n-r}b^{r} + \dots + b^{n},$$

where *n* is a positive integer and
$$\binom{n}{r} = \frac{n!}{(n-r)! r!} = \frac{n(n-1)\dots(n-r+1)}{r!}$$

2. TRIGONOMETRY

Identities

$$\sin^2 A + \cos^2 A = 1.$$

$$\sec^2 A = 1 + \tan^2 A.$$

$$\csc^2 A = 1 + \cot^2 A.$$

$$\sin(A \pm B) = \sin A \cos B \pm \cos A \sin B$$

$$\cos(A \pm B) = \cos A \cos B \mp \sin A \sin B$$

$$\tan(A \pm B) = \frac{\tan A \pm \tan B}{1 \mp \tan A \tan B}$$

$$\sin 2A = 2 \sin A \cos A$$

$$\cos 2A = \cos^2 A - \sin^2 A = 2 \cos^2 A - 1 = 1 - 2 \sin^2 A$$

$$\tan 2A = \frac{2 \tan A}{1 - \tan^2 A}$$

Formulae for $\triangle ABC$

$$\frac{a}{\sin A} = \frac{b}{\sin B} = \frac{c}{\sin C}$$
$$a^{2} = b^{2} + c^{2} - 2bc \cos A$$
$$\Delta = \frac{1}{2} ab \sin C$$

1	(i) Differentiate	$\ln(\cos x)$ with respect to <i>x</i> .	[1]
	(ii) Hence find	$\int \tan x \mathrm{d}x .$	[2]

(iii) Differentiate $x \tan x$ with respect to x. [2]

(iv) Using the results from (ii) and (iii), hence find	$\int x \sec^2 x \mathrm{d}x .$	[3]
--	-----------------------------------	-----

2 (i) Prove that
$$\frac{(\sin A + \cos A)(1 - \sin A \cos A)}{\sin^3 A} = 1 + \cot^3 A$$
 [4]

(ii) Hence solve the equation $(\sin A + \cos A)(1 - \sin A \cos A) = 2\sin^3 A$ for [3] $0^\circ \le A \le 180^\circ$. (i) The expression 3x³ + hx² + kx-4, where h and k are constants, has a factor of 3x-1 and leaves a remainder of -4 when divided by x+1. By forming 2 equations, of h and of k, show that the value of h = 11 and k = 8. [4]

(ii) Hence, express
$$\frac{-x^2+6x+9}{3x^3+hx^2+kx-4}$$
 as the sum of three partial fractions [7]

A particle, moving in a straight line, passes through a fixed point O with a speed of 4 m/s. The acceleration, $a \text{ m/s}^2$, of the particle, t s after passing through O, is given

by
$$a = -\frac{4}{15}e^{-\frac{t}{60}}$$
.

4

(i) Find the exact value of t when the particle is at instantaneous rest.

(ii) Find an expression in term of *t*, for the displacement, from *O*, of the particle *t* seconds after passing *O*.

[3]

[6]

(iii) Hence find the distance of the particle from *O* when it is at instantaneous rest. [1]

4

(iv) Show that the particle is again at *O* at some instant during the thirty-seventh second [2] after first passing through *O*.

(i) The equation $\log_3 x - \frac{1}{2} \log_{27} x = \log_2 8$ has the solution $x = 3^k$. Find the value of k.

[4]

5

(ii) Solve the equation $2\log_4(x-2) - \log_4(x+10) = \frac{1}{2}$. [4]

The roots of the quadratic equation $2x^2 = x + 3$ are α and β . (i) Find the value of $\alpha^2 + \beta^2$. [2]

6

(ii) Find the quadratic equation whose roots are $\frac{\beta}{\alpha^2}$ and $\frac{\alpha}{\beta^2}$. [5]

7 The points A(2, 1) and B(11, -2) lie on a circle.

(i) Find the equation of the perpendicular bisector of the chord *AB*. [4]

The line with equation $y = \frac{4}{3}x - 10$ is a normal to the circle. (ii) Hence, find the equation of the circle.

[5]

[1]

(iii) Find the coordinates of the point on the circle which is at the greatest distance from the *x*-axis.

11



8

The diagram shows part of the graph of y = |3x-7|+2. A horizontal line is drawn from *A* to intersect the graph of y = |3x-7|+2 at *B*. (i) Find the coordinates of *A*, *B* and *C*.

[3]

(ii) Find the set of values of *m* for which the graph of y = |3x-7|+2 and the line y = mx intersects at 2 points.

[2]

[3]

(iii) Solve the equation |3x-7|+2=4x-1

8

13

The resistance to motion, R newtons, of a plank towed through water and its

speed, V m/s is given by $R = AV^n$, where A and n are constants. The table shows the corresponding values of V and R.

· · · · · · · · · · · · · · · ·				
V	1.65	2.46	3.68	6.05
R	1.87	3.70	7.33	17.1

9

(ii) Use the graph to estimate the value of each of the constants A and n,[4] giving your answers correct to 1 decimal place.

[3]

(iii) By drawing a suitable line on your graph, solve the equation $AV^n = V^{1,2}$, giving [3] your answer correct to 1 decimal place.

⁽i) On the grid on page 15, draw the graph of ln R against ln V, using a scale of 4 cm for 0.5 unit on the ln V-axis and a scale of 5 cm to 1 unit on the ln R-axis.

+ +


The diagram show part of the curve $y = 3 - \frac{18}{2x+3}$. The normal to the curve at x = 3 intersects the y-axis at A. Find the exact area of the shaded region. [11]

Continuation of Working Space for Question 10.



11

A rectangular table *PQRS* is positioned at a corner of a room. Given that PQ = 1.6m PS = 0.5 m, angle $OQP = \theta$ and angle QOP = angle $RTQ = 90^{\circ}$, where θ varies. (i) Show that $OT = 1.6 \cos \theta + 0.5 \sin \theta$.

End of Paper

Answer Key

1a	1 2	7i	Since the product of 2 gradients is -1 .
	$x = -\frac{1}{2}, y = \frac{1}{2}$ or $x = 1, y = -2$		AB is perpendicular to BC.
h	a = 23, b = -13	ii	Proof
2	<i>k</i> < -5	iii	M = (h, 3h) C(6h, 3h)
3a	1	iv	h = 5
Ju	$\frac{1}{2}$	1,	
h	$\frac{2}{\sqrt{15}}$ 12	8i	dy = 5
	$\frac{6\sqrt{13}-12}{11}$	01	$\left \frac{dy}{dt}\right = \frac{3}{2}$
4.	11		$ dx _{x=-2} = 2$
41	52.8%	11	2y = 5x + 50
11	16	9i	$\frac{dy}{dt} = -3\cos^2 x \sin x + 3\cos^2 x$
			dx
			$\frac{d^2y}{dx^2} = -3\cos^3 x + 6\sin^2 x \cos x - 9\sin^3 x$
	As t gets very large $e^{-0.15t}$	ii	dx^2
111	approaches zero but will never	11	$\mathbf{A} = 0$
	reach zero. Therefore, the		
	percentage will not reach 100%.		
5i	Proof	10i	Proof
ii	<i>n</i> = 12	ii	$(0, 5, w^2)$
			$60 - \frac{12}{12}x^{-1}$
iii	$-\frac{11264}{r^5}$	iii	2.5cm ² /s
	27 *		
6i	Proof	11i	Least value $= -4$
		••	Greatest value = 2
11	$\left(-\frac{2}{2}, -\sqrt{85}\right)$	11	π
	(9, ,)		
iii	minimum point	iii	
			-2 0 2
		1V 12	4 solutions
		12	$\frac{dy}{dx} = e^{2x-3}(x+1)^2[5+2x]$
			5
			$x > -\frac{1}{2}, 2x + 5 > 0, (x + 1)^2$ and e^{2x-3}
			> 0
			$\frac{dy}{dy} > 0$ the curve is increasing
			dx = -0, the curve is increasing
1		1	

Answer Key

1(i)	$-\tan x$	8(i)	$A = (0,9), B = \left(\frac{7}{3}, 2\right), C = \left(\frac{14}{3}, 9\right)$
(ii)	$-\ln(\cos x) + c$	(ii)	$\frac{6}{7} < m < 3$
(iii)	$\tan x + x \sec^2 x$	(iii)	$x = -4$ (rejected) or $x = \frac{10}{7}$
(iv)	$x\tan x + \ln(\cos x) + c$	9(ii)	$n = \text{gradient} = 1.7, \ A = e^{-0.24} = 0.8$
2(ii)	45°	(iii)	1.6
3(i)	$h + 3k = 35, \ h - k = 3$	10	$\frac{33}{8} + 9 \ln 3$
(ii)	$\frac{2}{3x-1} - \frac{1}{x+2} + \frac{1}{(x+2)^2}$	11(ii)	61.6°
4(i)	$-60\ln\frac{3}{4}$		
(ii)	$s = -960e^{-\frac{t}{60}} - 12t + 960$		
(iii)	32.9 m		
5(i)	$\frac{18}{5}$		
(ii)	x = 8 or $x = -2$ (rejected)		
6(i)	$\frac{13}{4}$		
(ii)	$x^2 - \frac{19}{18}x - \frac{2}{3} = 0$		
7(i)	$y = 3x - \overline{20}$		
(ii)	$(x-6)^2 + (y+2)^2 = 25$		
(iii)	(6, -7)		



ZHONGHUA SECONDARY SCHOOL PRELIMINARY EXAMINATION 2020 SECONDARY 4E/5N

Candidate's Name	Class	Register Number
MARK SCHEME		

ADDITIONAL MATHEMATICS

4047/01

PAPER 1

15 September 2020 2 hours

Candidates answer on the Question Paper

READ THESE INSTRUCTIONS FIRST

Write your name, class and register number on all the work you hand in. Write in dark blue or black pen on both sides of the paper. You may use an HB pencil for any diagrams or graphs. Do not use staples, paper clips, glue or correction fluid.

Answer **all** the questions.

Give non-exact numerical answers correct to 3 significant figures, or 1 decimal place in the case of angles in degrees, unless a different level of accuracy is specified in the question.

The use of a scientific calculator is expected, where appropriate.

You are reminded of the need for clear presentation in your answers.

The number of marks is given in brackets [] at the end of each question or part question.

The total number of marks for this paper is 80.



Mathematical Formulae

1. ALGEBRA

Quadratic Equation

For the equation $ax^2 + bx + c = 0$,

$$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$

Binomial Theorem

$$(a+b)^{n} = a^{n} + \binom{n}{1}a^{n-1}b + \binom{n}{2}a^{n-2}b^{2} + \dots + \binom{n}{r}a^{n-r}b^{r} + \dots + b^{n},$$

where *n* is a positive integer and
$$\binom{n}{r} = \frac{n!}{(n-r)! r!} = \frac{n(n-1)\dots(n-r+1)}{r!}$$

2. TRIGONOMETRY

Identities

$$\sin^2 A + \cos^2 A = 1.$$

$$\sec^2 A = 1 + \tan^2 A.$$

$$\csc^2 A = 1 + \cot^2 A.$$

$$\sin(A \pm B) = \sin A \cos B \pm \cos A \sin B$$

$$\cos(A \pm B) = \cos A \cos B \mp \sin A \sin B$$

$$\tan(A \pm B) = \frac{\tan A \pm \tan B}{1 \mp \tan A \tan B}$$

$$\sin 2A = 2 \sin A \cos A$$

$$\cos 2A = \cos^2 A - \sin^2 A = 2\cos^2 A - 1 = 1 - 2\sin^2 A$$

$$\tan 2A = \frac{2 \tan A}{1 - \tan^2 A}$$

Formulae for $\triangle ABC$

$$\frac{a}{\sin A} = \frac{b}{\sin B} = \frac{c}{\sin C}$$
$$a^{2} = b^{2} + c^{2} - 2bc \cos A$$
$$\Delta = \frac{1}{2} ab \sin C$$

1	(a)	Find the values of <i>x</i> and <i>y</i> which satisfy the equations.		
		$3^x \times \sqrt{3^y} = 1$		
		$4^{x-4} \div 32^y = 16^{\frac{1}{x}}$	1	[5]
		$3^{x} \times \sqrt{3^{y}} = 1 \qquad 4^{x-4} \div 32^{y} = 16^{\frac{1}{x}}$ $3^{x} \times 3^{\frac{1}{2}^{y}} = 3^{0} \qquad 2^{2x-8} \div 2^{5y} = 2^{\frac{4}{x}}$ $3^{x+\frac{1}{2}^{y}} = 3^{0} \qquad 2^{2x-8-5y} = 2^{\frac{4}{x}}$	[M1] applying laws of indices correctly	
		$x + \frac{1}{2}y = 0 \qquad 2x - 8 - 5y = \frac{4}{x}$ $x = -\frac{1}{2}y - (1) \qquad 2x^2 - 5xy - 8x - 4 = 0 - (2)$	[A1] for either (1) or (2) correctly)
		Sub (1) into (2): $2\left(-\frac{1}{2}y\right)^{2} - 5\left(-\frac{1}{2}y\right)y - 8\left(-\frac{1}{2}y\right) - 4 = 0$ (1)	[M1] substitution	
		$2\left(\frac{1}{4}y^{2}\right) + \frac{5}{2}y^{2} + 4y - 4 = 0$ $3y^{2} + 4y - 4 = 0$ (3y - 2)(y + 2) = 0 $y = \frac{2}{3} \text{or} y = -2$		
		$y = \frac{2}{3}$ $x = -\frac{1}{2}\left(\frac{2}{3}\right) = -\frac{1}{3}$	[A1] both x and y corre	ct
		When $y = -2$ $x = -\frac{1}{2}(-2) = 1$	[A1] both x and y corre	ct

(b)	Without using a calculator, find the values of a and b such that		
	$7-3\sqrt{3} = (a+b\sqrt{3})(2+\sqrt{3})$, where a and b are integers.		[4]
	$7 - 3\sqrt{3} = (a + b\sqrt{3})(2 + \sqrt{3})$ $a + b\sqrt{3} = \frac{7 - 3\sqrt{3}}{2 + \sqrt{3}}$ $= \frac{7 - 3\sqrt{3}}{2 + \sqrt{3}} \times \frac{2 - \sqrt{3}}{2 - \sqrt{3}}$ $7(2 - \sqrt{3}) - 3\sqrt{3}(2 - \sqrt{3})$	[M1] rationalisation	
	$= \frac{14 - 7\sqrt{3} - 6\sqrt{3} + 9}{4 - 3}$ = 23 - 13 \sqrt{3}	[M1] expansion/simplification	on
	$a = 23, \qquad b = -13$ Alternative solution	[A1] [A1]	
	$a(2+\sqrt{3})+b\sqrt{3}(2+\sqrt{3})=7-3\sqrt{3}$ $2a+a\sqrt{3}+2b\sqrt{3}+3b=7-3\sqrt{3}$ $2a+3b+a\sqrt{3}+2b\sqrt{3}=7-3\sqrt{3}$		
	2a + 3b = 7 (1) a + 2b = -3 (2) (2) into (1): 2(-3 - 2b) + 3b = 7 -6 - 4b + 3b = 7 b = -13 a = 23	[M1] grouping[M1] simultaneous equation[A1][A1]	

2	Find the set of values of the constant k for which the curve	$w = (k+2)x^2 - 10x + 2k + 1$ lies
	completely below the line $y = 2x + 3$.	[4]
	$(k+2)x^{2} - 10x + 2k + 1 < 2x + 3$ (k+2)x ² - 10x + 2k + 1 - 2x - 3 < 0 (k+2)x ² - 12x + 2k - 2 < 0	[M1] eliminate y
	$b^{2} - 4ac < 0 \qquad \text{and} \qquad k + 2 < 0$ $(-12)^{2} - 4(k+2)(2k-2) < 0 \qquad \text{and} \qquad k < -2$ $144 - (8k^{2} + 8k - 16) < 0$ $-8k^{2} - 8k + 160 < 0$ $k^{2} + k - 20 > 0$	[B1] use of correct discriminant
	(k-4)(k+5) > 0 k < -5 or $k > 4\therefore k < -5$	[A1] [A1]

3	Given that $\sin A = \frac{2}{3}$ and $\tan B = -\frac{1}{\sqrt{3}}$ where angles A and B lie in the same quadrant,				
	leavi	leaving your answers in exact form, calculate the value of			
	(a)	$\cos 2B$			
		$\cos 2B = 2\cos^2 B - 1$ $= 2\left(-\frac{\sqrt{3}}{2}\right)^2 - 1$ $= 2\left(\frac{3}{4}\right) - 1$ $= \frac{1}{2}$	[M1] Applying double angle [A1]		
	(b)	$\sec(A-B)$			
		$\sec(A-B) = \frac{1}{\cos(A-B)}$ $= \frac{1}{\cos A \cos B + \sin A \sin B}$ $= 1 \div \left(-\frac{\sqrt{5}}{3} \times -\frac{\sqrt{3}}{2} + \frac{2}{3} \times \frac{1}{2}\right)$ $= 1 \div \left(\frac{\sqrt{15}}{6} + \frac{1}{3}\right)$ $= 1 \div \left(\frac{\sqrt{15}+2}{6}\right)$ $= \frac{6}{\sqrt{15}+2}$ $= \frac{6\sqrt{15}-12}{11}$	[B1] change from sec to cos [M1] addition formula		

4	In view of a contagious virus, the government of a particular country has imposed a 'Stay-Home-Notice' on her people to reduce the number of human-to-human transmission cases. It is estimated that the percentage of the population, <i>P</i> , complying to the Stay-Home-Notice is given by the equation $P = 100(1 - e^{-0.15t})$, where <i>t</i> is the number of days after the imposition			
	(i)	Find the percentage of population complying to the 'Stay-H	Iome-Notice' after 5	[1]
		days of the imposition. $P = 100(1 - e^{-0.15t})$ $= 100(1 - e^{-0.15\times5})$ $= 52.763$ $= 52.8\%$	[B1]	
	(ii)	Find the number of days after the imposition that it will tak the population to comply.	te for at least 90% of	[2]
		90 = 100 $(1 - e^{-0.15t})$ 0.9 = 1 - $e^{-0.15t}$ 0.1 = $e^{-0.15t}$ ln $e^{-0.15t}$ = ln 0.1 -0.15t = ln 0.1 t = 15.35 ≈ 15.4 No. of complete days = 16	[M1] take ln [A1]	
	(iii)Is it possible for the percentage of this country's population complying to the 'Stay-Home-Notice' to reach 100%? Explain your answer.			[1]
		As t gets very large, $e^{-0.15t}$ approaches zero but will never reach zero. Therefore, the percentage will not reach 100%.	[B1]	

5	(i)	In the expansion of $\left(2-\frac{x}{3}\right)^n$, show that the ratio of coefficient of the 2 nd term		
		to that of the 4 th torm can be simplified to the expression -	216	[4]
		to that of the 4 term can be simplified to the expression ((n-1)(n-2).	
		$\left(2-\frac{x}{3}\right)^n$		
		$=2^{n}-\binom{n}{1}2^{n-1}\binom{x}{3}+\binom{n}{2}2^{n-2}\binom{x}{3}^{2}-\binom{n}{3}2^{n-3}\binom{x}{3}^{3}+\dots$	[M1] binomial theore general term used	m or
		$=2^{n}-\frac{2^{n-1}nx}{3}+\frac{2^{n-2}n(n-1)x^{2}}{2\times 9}-\frac{2^{n-3}n(n-1)(n-2)x^{3}}{6\times 27}+\dots$		
		$-\frac{2^{n-1}n}{3}$	[B1]	
		Cofficient of $\frac{1}{T_4} = \frac{1}{2^{n-3}n(n-1)(n-2)}$	[B1]	
		$=\frac{54 \times 2^{n-1-n+3}}{(n-1)(n-2)}$	[M1] simplification leading to correct ans	wer
		$=\frac{54 \times 2^2}{(n-1)(n-2)}$		
		$=\frac{216}{(n-1)(n-2)}$	A.G.	

(ii)	Find the value of n if the ratio in (i) is $108:55$.	[2]
(ii)	Find the value of <i>n</i> if the ratio in (i) is 108:55. $\frac{216}{(n-1)(n-2)} = \frac{108}{55}$ 11880 = 108 ($n^2 - 3n + 2$) 110 = $n^2 - 3n + 2$ $n^2 - 3n - 108 = 0$ ($n - 12$)($n + 9$) = 0 n = 12 or $n = -9$ (rejected)	[M1] [A1]
(iii)	Hence, find the term in x^5 .	[2]
	$T_{6} = {\binom{12}{5}} 2^{12-5} {\binom{-x}{3}}^{5}$ $= -\frac{11264}{27} x^{5}$	[M1] [A1]

6	The equation of a curve is $y = \frac{2x-9}{\sqrt{x^2+1}}$.			
	(i)	Show that $\frac{dy}{dx} = \frac{9x+2}{\sqrt{(x^2+1)^3}}$.		[3]
		$y = \frac{2x-9}{\sqrt{x^2+1}}$ $\frac{dy}{dx} = \frac{\left(x^2+1\right)^{\frac{1}{2}}(2) - (2x-9)\frac{1}{2}\left(x^2+1\right)^{-\frac{1}{2}}(2x)}{x^2+1}$ $= \frac{2\left(x^2+1\right)^{\frac{1}{2}} - x(2x-9)\left(x^2+1\right)^{-\frac{1}{2}}}{x^2+1}$	[M1] quotient rule [B1] correct expressi	on
		$=\frac{\left(x^{2}+1\right)^{-\frac{1}{2}}\left[2\left(x^{2}+1\right)-x(2x-9)\right]}{x^{2}+1}$ $=\frac{\left(x^{2}+1\right)^{-\frac{1}{2}}\left[2x^{2}+2-2x^{2}+9x\right]}{x^{2}+1}$	[A1] factorisation lea to correct answer	nding
		$=\frac{2+9x}{\left(x^{2}+1\right)^{\frac{3}{2}}}$	[A.G]	

(ii)	Find the coordinates of the stationary point of the curve, leaving your answer in exact form.		[3]
	For stationery points.		1
	$\frac{dy}{dx} = 0$	[M1]	
	$\frac{2+9x}{x^3} = 0$		
	$(x^2+1)^2$ 2+9x=0		
	$x = -\frac{2}{9}$	[A1]	
	$\therefore y = \frac{2\left(-\frac{2}{9}\right) - 9}{\sqrt{\left(-\frac{2}{9}\right)^2 + 1}}$		
	$=\frac{-\frac{4}{9}-9}{\sqrt{\frac{85}{81}}}$		
	$= \frac{-\frac{85}{9}}{\frac{\sqrt{85}}{9}}$ $= -\frac{-\frac{85}{\sqrt{85}}}{\sqrt{85}} = -\sqrt{85}$		
	Coordinates = $\left(-\frac{2}{9}, -\sqrt{85}\right)$	[A1]	
 (iii)	Find the nature of this stationary point.		[2]
	$\begin{array}{c c} \mathbf{x} & \left(-\frac{2}{9}\right)^{-} & -\frac{2}{9} & \left(-\frac{2}{9}\right)^{+} \\ \hline \text{Sign} & - & 0 & + \\ \hline \text{slope} & & & & \\ \end{array}$	[M1]	
	$\left(-\frac{2}{9}, -\sqrt{85}\right)$ is a minimum point.	[A1]	



(iii)	Express the coordinates of <i>M</i> and of <i>C</i> in terms of <i>h</i> .	[3]
	$A(0,h)$ $M = \left(\frac{2h+0}{2}, \frac{5h+h}{2}\right)$ $= (h,3h)$ When $y = 3h$, x + 2(3h) = 12h x + 6h = 12h x = 6h C(6h,3h)	[B1] [M1] [A1]
(iv)	Find the value of h if the area of triangle AMC is 125 squar Area $= \frac{1}{2} \begin{vmatrix} 0 & 6h & h & 0 \\ h & 3h & 3h & h \end{vmatrix}$ $= \frac{1}{2} (18h^2 + h^2 - 6h^2 - 3h^2)$ $= 5h^2$ $\therefore 5h^2 = 125$ $h^2 = 25$ $h = \sqrt{25}$ $= 5$	[M1]

8	The equation of a curve is $y = \frac{x-6}{x+4} + 4$, where $x \neq -4$.				
	Find the				
	(i) gradient of the tangent at the point where the curve passes through the <i>x</i> -axis,				
		When $y = 0$ $\frac{x-6}{x+4} + 4 = 0$ $x-6 = -4x - 16$ $5x = -10$	[M1]		
		$x = -2$ $\frac{dy}{dx} = \frac{x + 4 - (x - 6)}{(x + 4)^2}$ $= \frac{10}{(x + 4)^2}$	[M1] [A1]		
		$\frac{dy}{dx}\Big _{x=-2} = \frac{10}{(-2+4)^2}$ $= \frac{5}{2}$	[A1]		

(ii)	equation of another tangent to the curve that is parallel to the tangent in (i). [4]			
	$\frac{dy}{dx} = \frac{5}{2}$	[M1]		
	$\frac{10}{(1-1)^2} = \frac{5}{2}$			
	$(x+4)^2 = 2$ $(x+4)^2 = 4$			
	$x + 4 = \pm 2$			
	x = -2 or $x = -6$	[A1]		
	$x = -6$, $y = \frac{-6-6}{-6+4} + 4 = 10$			
	Equation:			
	$\frac{y-10}{x+6} = \frac{5}{2}$	[M1]		
	2y - 20 = 5x + 30	[A1]		
	2y = 5x + 50	[]		

9
 The equation of a curve is
$$y = \cos^3 x + \sin 3x$$
.
 [4]

 (i)
 Find expressions for $\frac{dy}{dx}$ and $\frac{d^2y}{dx^2}$.
 [4]

 y = $\cos^3 x + \sin 3x$
 [4]

 $\frac{dy}{dx} = 3\cos^2 x(-\sin x) + 3\cos 3x$
 [M1] chain rule seen

 $\frac{dy}{dx} = -3\cos^2 x \sin x + 3\cos 3x$
 [A1]

 $\frac{d^2y}{dx^2} = -3[\cos^2 x \cos x + \sin x(2\cos x)(-\sin x)] - 9\sin 3x$
 [M1] product rule seen

 $= -3\cos^3 x + 6\sin^2 x \cos x - 9\sin 3x$
 [A1]

(ii)	Given that $\frac{d^2 y}{dx^2} + 9y = A\cos x$, find the value of A.	[2]		
$-3\cos^{3}x + 6\sin^{2}x\cos x - 9\sin 3x + 9\cos^{3}x + 9\sin 3x$ = $6\cos^{3}x + 6\sin^{2}x\cos x$ = $6\cos x(\cos^{2}x + \sin^{2}x)$ [M1] identity seen = $6\cos x(1)$				
A = 6	5 [A1]			



(iii)	Find the rate of change of the shaded area when PQ is halfway towards AB from C .	[3]
	$\frac{dA}{dt} = \frac{dA}{dx} \times \frac{dx}{dt} \qquad [B1] use \ of \ chain \ rule \ correctly$ $\frac{dA}{dx} = -\frac{10}{12}x \qquad [B1]$	
	$\frac{dA}{dt} = -\frac{10}{12}(6) \times (0.5)$	
	$= -2.5 \qquad [B1]$ Area is decreasing at a rate of 2.5cm ² /s	



(iv)	State, with detailed workings, the number of solutions of the equation	
	$3\pi\cos 2x = 2x - \pi$ for $-\pi \le x \le \pi$.	[2]

 $3\pi \cos 2x = 2x - \pi$ $3\cos 2x = \frac{2x}{\pi} - 1$ $3\cos 2x - 1 = \frac{2x}{\pi} - 2$ [B1] for manipulation

Since there are 4 intersection points, there are 4 solutions [B1] dep

The function f is defined by $y = (x+1)^3 e^{2x-3}$, for $x > -\frac{5}{2}$, and $x \neq -1$. Explain, with 12 [4] working, whether f is an increasing or decreasing function.

$$\frac{dy}{dx} = 3(x+1)^2 e^{2x-3} + 2(x+1)^3 e^{2x-3} \qquad [M1] \text{ product rule } [A1]$$
$$\frac{dy}{dx} = e^{2x-3}(x+1)^2 [3+2(x+1)]$$
$$\frac{dy}{dx} = e^{2x-3}(x+1)^2 [5+2x]$$

Given that

$$x > -\frac{5}{2}$$
, $2x + 5 > 0$, $(x + 1)^2$ and $e^{2x-3} > 0$, [M1]

 $\therefore \frac{dy}{dx} > 0$, the curve is increasing [A1] dep.

1 (i) Differentiate $\ln(\cos x)$ with respect to x.					
Answer	1	Marks	Guidance		
$\frac{d}{dr}\ln(co)$	$sx) = \frac{1}{\cos x} \times (-\sin x)$				
dx	$\frac{\cos x}{\sin x}$	h			
	$= \frac{-\frac{1}{\cos x}}{\cos x}$	– B1			
	$=-\tan x$				
1	(ii) Hence find $\int \tan x dx$			[2]	
		I			
Answer		Marks	Guidance		
$\int \tan x d$	x				
$=-\int -t$	an $x dx$	M1	Reverse differentiation		
$=-\ln(cc)$	$(\cos x) + c$	A1			
1	(iii) Differentiate wten w with respect to w			[2]	
1	(iii) Differentiate x tai x with respect to x				
Answer		Marks	Guidance		
$d(x \tan x) = \tan x + x \cos^2 x$			Product rule		
$\frac{dx}{dx}$	$(x) = \tan x + x \sec x$	A 1			
		AI			
1	(iv) Using the results from (ii) and (iii), he	ence find	$\int x \sec^2 x \mathrm{d}x$	[3]	
		1		·	
Answer		Marks	Guidance		
$x \tan x$ -	$+c_1 = \int \tan x + x \sec^2 x \mathrm{d}x$	M1	Reverse differentiation		
$x \tan x + \mathbf{c}_1 - \int \tan x \mathrm{d}x = \int x \sec^2 x \mathrm{d}x$			Split to 2 integrals		
$\int x \sec^2 x$	$x \mathbf{d}x = x \tan x - (-\ln(\cos x)) + \mathbf{c}$				
$= x \tan x + \ln(\cos x) + c \qquad A1$					

2 (i) Prove that
$$\frac{(\sin A + \cos A)(1 - \sin A \cos A)}{\sin^3 A} = 1 + \cot^3 A$$
 [4]

Answer	Marks	Guidance
$\sin A + \cos A - \sin^2 A \cos A - \sin A \cos^2 A$	M1	Correct expansion
$LHS sin^3 A$	D1	
$\sin A + \cos A - (1 - \cos^2 A) \cos A - \sin A(1 - \sin^2 A)$	BI	Use of $2 4 \cdot 1 = 2 4 \cdot 1$
$\equiv \frac{1}{\sin^3 A}$		$\cos^2 A + \sin^2 A = 1$
$= \sin A + \cos A - \cos A + \cos^3 A - \sin A + \sin^3 A$	B1	
$-\frac{1}{\sin^3 A}$		
$=$ $\frac{\cos^3 A + \sin^3 A}{\sin^3 A}$		
$\sin^3 A$		
$=$ $\frac{\cos^3 A}{\cos^3 A}$ $\pm \frac{\sin^3 A}{\cos^3 A}$	B1	
$\sin^3 A \sin^3 A$		
$= 1 + \cot^3 A$		

Alternative solution

Answer	Marks	Guidance		
$\lim_{HS^{-}} (\sin A + \cos A) \qquad (1 - \sin A \cos A)$	B1			
$\lim_{n \to \infty} \frac{1}{\sin A} \times \frac{1}{\sin^2 A}$	DI			
$= (1 + \cot A) \times (1 + \sin A \cos A)$	BI	Use of $\frac{\cos A}{\sin A}$		
$=$ (1 + cot A) × $\left(\frac{1}{\sin^2 A} - \frac{1}{\sin^2 A}\right)$		sin A		
$= (1 + \cot A) \times (\cos ec^2 A - \cot A)$				
	B1	Use of		
$= (1 + \cot A) \times (1 + \cot^2 A - \cot A)$		$1 + \cot^2 A = \cos ec^2 A$		
= $1 + \cot A - \cot^2 A + \cot^3 A - \cot A - \cot^2 A$	D1			
= 1 + cot ³ A	BI	Correct expansion		

2	(ii) Hence solve the equation	$(\sin A + \cos A)(1 - \sin A \cos A) = 2\sin^3 A$ for	[3]
	$0^{\circ} \le A \le 180^{\circ}.$		

Answer	Marks	Guidance
$\frac{(\sin A + \cos A)(1 - \sin A \cos A)}{\sin^3 A} = 2$ $1 + \cot^3 A = 2$	B1	
$\cot^{3}A = 1$ $\cot A = 1$ $\tan A = 1$	M1	Taking cube root without negative sign
$A = 45^{\circ}$	A1	

3	(i) The expression $3x^3 + hx^2 + kx - 4$, where h and k are constants, has a	
	factor of $3x-1$ and leaves a remainder of -4 when divided by $x+1$.	
	By forming 2 equations of <i>h</i> and of <i>k</i> , show that the value of $h = 11$ and $k = 8$.	[4]
	By forming 2 equations of <i>n</i> and of <i>k</i> , show that the value of <i>n</i> . If and <i>k</i> = 0.	[']

Answer	Marks	Guidance
Let $f(x) = 3x^3 + hx^2 + kx - 4$		
(3x-1) is a factor of f (x)		(1)
$f\left(\frac{1}{3}\right) = 0$	M1	Realise $f\left(\frac{1}{3}\right) = 0$
$3 \times \left(\frac{1}{3}\right)^3 + h\left(\frac{1}{3}\right)^2 + k\left(\frac{1}{3}\right) - 4 = 0$		
$\frac{1}{9} + \frac{h}{9} + \frac{k}{3} - 4 = 0$		
h + 3k = 35(1)		
Divisor = $x + 1$, reminder = -4		
f(-1) = -4		
-3 + h - k - 4 = -4		
h - k = 3(2)	M1	Realise $f(-1) = -4$
(1) - (2), 4k = 32		
k = 8	DI	DM on correct equation (1)
h = 3 + k	DMI B1	and (2)
=3+8=11	DI	

|--|

Answer	Marks	Guidance
Let $3x^3 + 11x^2 + 8x - 4 = (3x - 1)(x^2 + bx + 4)$		
comparing the coefficient of x^2 ,		
11 = -1 + 3b		
b = 4		
$3x^{3} + 11x^{2} + 8x - 4 = (3x - 1)(x^{2} + 4x + 4)$	M1	Long division or inspection
$= (3x-1)(x+2)^2$	A1	
$-x^2+6x+9$ A B C		
$\frac{1}{(3x-1)(x+2)^2} = \frac{1}{3x-1} + \frac{1}{x+2} + \frac{1}{(x+2)^2}$	M1	
$-x^{2} + 6x + 9 = A(x+2)^{2} + B(3x-1)(x+2) + C(3x-1)$	M1	
Let $x = -2$		
$-(-2)^{2}+6(-2)+9=C(-7)$		
-4 - 12 + 9 = -7C		
C = 1		
Let $x = \frac{1}{3}$		
$-\frac{1}{9} + \frac{6}{3} + 9 = A\left(\frac{7}{3}\right)^2$		
98 49 <i>A</i>		
$\frac{1}{9} = \frac{1}{9}$		
A = 2		
comparing coefficients of x^2 ,		
-1 = A + 3B		A1 for each correct A, B, C
-1 - 2 = 3B		
B = -1		
$\frac{-x^2 + 6x + 9}{-x^2 + 6x + 9} = \frac{2}{-x^2 - 1} + \frac{1}{-x^2 - 1}$		
$(3x-1)(x+2)^2 = 3x-1 + x+2 + (x+2)^2$		

4	A particle, moving in a straight line, passes through a fixed point O with a speed of 4 m/s. The acceleration, $a \text{ m/s}^2$, of the particle, t s after passing through O, is given by $a = -\frac{4}{15}e^{-\frac{t}{60}}$.	
	(i) Find the exact value of t when the particle is at instantaneous rest.	[6]

Answer	Marks	Guidance
$v = \int -\frac{4}{15} e^{-\frac{t}{60}} dx$	M1	Integrating <i>a</i>
$= -\frac{4}{15} \times \frac{e^{-\frac{t}{60}}}{-\frac{1}{60}} + c$	A1	Correct integration (accept without + +c)
$v = 16e^{-\frac{1}{60}} + c$		
when $t = 0$, $v = 4$		
$4 = 16e^0 + c$		
c = -12		
$v = 16e^{-\frac{t}{60}} - 12$	A1	
At instantaneous rest, $v = 0$		
$16e^{-\frac{t}{60}} - 12 = 0$	M1	Equating <i>v</i> to zero
$e^{-\frac{t}{60}} = \frac{12}{16}$		
$-\frac{t}{60} = \ln\frac{3}{4}$	M1	Taking logarithm
$t = -60\ln\frac{3}{4}$	A1	

4	(ii) Find an expression in term of t, for the displacement, from O, of the	
	particle <i>t</i> seconds after passing O.	[3]

Answer	Marks	Guidance
$s = \int 16e^{-\frac{t}{60}} - 12 \mathrm{d}t$	M1	Integrating to get s
$= \frac{16e^{-\frac{t}{60}}}{-\frac{1}{60}} - 12t + c$ = $-960e^{-\frac{t}{60}} - 12t + c$ when $t = 0, s = 0$ $0 = -960e^{0} - 0 + c$	A1	
960 = c		
$s = -960e^{-\frac{t}{60}} - 12t + 960$	A1	

(ii) Hence find the distance of the particle from O when it is at instantaneous	[1]
rest.	

Answer	Marks	Guidance
when $v = 0, t = -60 \ln \frac{3}{4}$		
$s = -960 \times \frac{3}{4} + 720 \ln \frac{3}{4} + 960$ = 32.9 m	A1	

(iv) Show that the particle is again at O at some instant during the	[2]
thirty-sixth second after first passing through O.	

Answer	Marks	Guidance
when $t = 36$, $s = -960e^{-\frac{36}{60}} - 12(36) + 960$		
= 1.14 m		
when $t = 37$, $s = -960e^{-\frac{37}{60}} - 12(37) + 960$	B1	
= -2.15 m When t=36, particle is on the right side of O with displacement of 1.14 and when t=37, particle is on the left side of O with s=-2.15. Hence at some instant during the 37 th second the particle is again at O.	B1	

5	(i) The equation $\log_3 x - \frac{1}{2}\log_{27} x = \log_2 8$ has the solution $x = 3^k$.	
	Find the value of k.	[4]

Answer	Marks	Guidance
$\log_3 x - \frac{1}{2} \times \frac{\log_3 x}{\log_3 27} = \log_2 2^3$	M1	Change of base law
$\log_3 x - \frac{1}{6}\log_3 x = 3$		
$\frac{5}{6}\log_3 x = 3$	M1	Simplify to single log
$\log_3 x = \frac{18}{5}$		
$x = 3^{\frac{18}{5}}$	A1	
By comparing with 3^k		
$k = \frac{18}{5}$ or $k = 3.6$	A1	

Alternatively

Answer	Marks	Guidance
since $x = 3^k$ is a solution,		
$\log_3 3^k - \frac{1}{2} \times \log_{27} 3^k = \log_2 2^3$	M1	Substitution
$k \log_3 3 - \frac{1}{2} \times \frac{\log_3 3^k}{\log_3 27} = 3$	M1	Applying power law
$k - \frac{1}{2} \times \frac{k \log_3 3}{\log_3 3^3} = 3$		
$k - \frac{1}{2} \times \frac{k \log_3 3}{3 \log_3 3} = 3$		
$k - \frac{1}{2} \times \frac{k}{3} = 3$		
$\frac{5k}{6} = 3$	B1	Reduce to a linear equation
$k = \frac{18}{5} = 3.6$	A1	

5 (ii) Solve the equation
$$2\log_4(x-2) - \log_4(x+10) = \frac{1}{2}$$
. [4]

Answer	Marks	Guidance
$2\log_4(x-2) - \log_4(x+10) = \frac{1}{2}$		
$\log_4 \frac{(x-2)^2}{(x+10)} = \frac{1}{2}$	M1	Apply power or subtraction law
$\frac{(x-2)^2}{(x+10)} = 4^{\frac{1}{2}}$	M1	convert to index form or equivalent.
$\left(x-2\right)^2 = 2\left(x+10\right)$		
$x^2 - 4x + 4 - 2x - 20 = 0$		
$x^2 - 6x - 16 = 0$		
(x-8)(x+2) = 0		
x = 8 or $x = -2$ (rejected)	A1 A1	x = -2, must be rejected

6	The roots of the quadratic equation $2x^2 = x + 3$ are α and β .	
	(i) Find the value of $\alpha^2 + \beta^2$.	[4]

Answer	Marks	Guidance
$2x^{2} = x + 3$ $\alpha + \beta = -\left(\frac{-1}{2}\right) = \frac{1}{2}$ $\alpha\beta = -\frac{3}{2}$ $(\alpha + \beta)^{2} = \alpha^{2} + 2\alpha\beta + \beta^{2}$ $\alpha^{2} + \beta^{2} = (\alpha + \beta)^{2} - 2\alpha\beta$	M1	
$= \left(\frac{1}{2}\right)^2 - 2 \times \left(-\frac{3}{2}\right)$ $= \frac{13}{4}$	A1	

6	(ii) Find the quadratic equations whose roots are $\frac{\beta}{\alpha^2}$ and $\frac{\alpha}{\beta^2}$.	[3]
---	---	-----
Answer	Marks	Guidance
--	-------	---------------
sum of roots = $\frac{\beta}{\alpha^2} + \frac{\alpha}{\beta^2}$		
$=\frac{\beta^3+\alpha^3}{\alpha^2\beta^2}$	M1	
$=rac{ig(lpha+etaig)ig(lpha^2-lphaeta+eta^2ig)}{lpha^2eta^2}$	B1	or equivalent
$=\frac{\left(\frac{1}{2}\right)\left(\frac{13}{4}-\left(-\frac{3}{2}\right)\right)}{2}$		
$\frac{9}{4}$		
$= \frac{1}{2} \times \frac{19}{4} \times \frac{4}{9}$		
$=\frac{19}{18}$	A1	
product of roots = $\frac{\beta}{\alpha^2} \times \frac{\alpha}{\beta^2} = \frac{1}{\alpha\beta} = -\frac{2}{3}$	A1	
Quadratic equation is $x^2 - \frac{19}{18}x - \frac{2}{3} = 0$	A1	

7	The points A(2,1) and B(11, -2) lie on a circle.	
	(i) Find the equation of the perpendicular bisector of the chord AB.	[4]

Answer	Marks	Guidance
gradient of $AB = \frac{3}{-9} = -\frac{1}{3}$		
gradient of the perpendicular bisector = $-\frac{1}{-\frac{1}{2}}$	M1A1	
= 3		
midpoint of $AB = \left(\frac{13}{2}, -\frac{1}{2}\right)$	B1	
Equation of perpendicular bisector is		
$y + \frac{1}{2} = 3\left(x - \frac{13}{2}\right)$	Al	
y = 3x - 20		

7	The line with equation $y = \frac{4}{3}x - 10$ is a normal to the circle.	
	(ii) Hence, find the equation of the circle.	[5]

Answer	Marks	Guidance
For centre of circle, $\frac{4}{3}x - 10 = 3x - 20$ $\frac{5}{3}x = 10$ $x = 6$	M1A1	Equating eqn of normal to eqn of perpendicular bisector A1 for the value of <i>x</i> .
y = 18 - 20 = -2 centre = (6, -2)	B1	
radius, $r = \sqrt{(6-2)^2 + (-2-1)^2}$ or $r = 11-6$ = 5 = 5	A1	
Equation of circle is $(x-6)^2 + (y+2)^2 = 25$	Al	

7 (iii) Find the coordinates of the point on the circle which is at the greatest distance [1] from the *x*-axis.

Answer	Marks	Guidance
point = (6, -7)	A1	



Answer	Marks	Guidance
At <i>A</i> , $y = 0, x = 9$		
A = (0,9)	A1	
At $C, y = 9$,		
9 = 3x - 7 + 2		
3x-7 = 7		
3x - 7 = 7		
$x = \frac{14}{3}$		
$C = \left(\frac{14}{3}, 9\right)$	A1	
$B = \left(\frac{7}{3}, 2\right)$	A1	

8	(ii) Find the set of values of <i>m</i> for which the graph of $y = 3x-7 +2$ and the line	[2]
	y = mx intersects at 2 points.	

[3]

Answer	Marks	Guidance
gradient $OB = \frac{2}{\frac{7}{3}} = \frac{6}{7}$	M1	
$\frac{6}{7} < m < 3$	A1	

	(iii)	Solve the equation	3x-7 +2=4x-1
--	-------	--------------------	--------------

8

Answer		Marks	Guidance
3 <i>x</i>	x-7 +2=4x-1		
3x	x-7 = 4x - 3		
3 <i>x</i>	$z - 7 = \pm (4x - 3)$	M1	
3 <i>x</i>	x - 7 = 4x - 3 or $3x - 7 = -(4x - 3)$		
<i>x</i> =	=-4(rejected) or	A1	Must be rejected
7 <i>x</i>	c = 10		
<i>x</i> =	$=\frac{10}{7}$	A1	

9	The resistance to motion, <i>R</i> newtons, of a plank towed through water and its							
	speed, V m/s is given by $R = AV^n$, where A and n are constants.							
	The table sl	nows the corr	responding v	alues of V an	d <i>R</i> .			
	V 1.65 2.46 3.68 6.05							
	R	1.87	3.70	7.33	17.1			
	(i) On the grid on page 15, draw the graph of ln <i>R</i> against ln <i>V</i> , using a						[3]	
	scale of 4 cm for 0.5 unit on the ln V-axis and a scale of							
	5 cm t	o 1 unit on th	e ln <i>R</i> -axis.					

Answer			Marks	Guidance		
$\ln R = \ln A + n \ln V$						
ln V	0.501	0.900	1.30	1.80		
ln <i>R</i>	0.626	1.31	1.99	2.84		
Axes and scale A1 All points plotted correctly join with a straight line				A2	Deduct 1 mark for any point plotted wrongly	

(ii) Use the graph to estimate the value of each of the constants A and n, giving	[4]
your answers to 1 decimal place.	

Answer		Marks	Guidance
	From the graph, $\ln r$ -intercept = -0.24	M1 A1	
	$\ln A = \ln r$ -intercept = -0.24		
	$A = e^{-0.24} = 0.8$ (to dec pl)		
1	gradient = $\frac{2.84 - (-0.24)}{1.80 - 0} = 1.7$	M1	Must indicate the 2 points
	n = gradient = 1.7 (to 1 dec pl)	A1	used to find gradient

(iii) By drawing a suitable line on your graph, solve the equation $AV^n = V^{1,2}$, giving	[4]
your answer correct to 1 decimal place.	

Answer	Marks	Guidance
$\ln A + n \ln V = 1.2 \ln V$		
Draw $\ln R = 1.2 \ln V$	B1	Correct line drawn
From the graph, $\ln V = 0.475$	M1	Reading off the lnV
$V = e^{0.475}$		coordinate from the point of
= 1.6 (to 1 dec pl)	A1	intersection



11	A rectangular table <i>PQRS</i> is positioned at a corner of a room. Given that $PQ = 1.6m$	
	$PS = 0.5$ m, angle $OQP = \theta$ and angle $QOP =$ angle $RTQ = 90^{\circ}$, where θ varies.	
	(i) Show that $OT = 1.6 \cos \theta + 0.5 \sin \theta$.	[3]
	P O O O O O O O O O O	

Answer	Marks	Guidance
$\angle RQT = 180^{\circ} - 90^{\circ} - \theta = 90^{\circ} - \theta$		$\angle QRT = \theta$ can be in the
$\angle QRT = 180^{\circ} - (90^{\circ} + 90^{\circ} - \theta) = \theta$	M1	diagram
$\cos\theta = \frac{OQ}{1.6}$	B1	
$OQ = 1.6\cos\theta$		
$\sin\theta = \frac{QT}{0.5}$	B1	
$QT = 0.5\sin\theta$		
OT = OQ + QT		Award 3 marks only if it is
$= 1.6\cos\theta + 0.5\sin\theta$		complete.

11 (ii) Find the value of θ for which $OT = 1.2$ m			
Answer	Marks	Guidance	
Let $1.6\cos\theta + 0.5\sin\theta = R\cos(\theta - \alpha)$	M1		
$R = \sqrt{1.6^2 + 0.5^2} = \sqrt{2.81}$	B1		
$\alpha = \tan^{-1} \frac{0.5}{1.6} = 17.354^{\circ}$	B1		
$\sqrt{2.81}\cos(\theta - 17.354^\circ) = 1.2$			
$\cos(\theta - 17.354^{\circ}) = \frac{1.2}{\sqrt{2.81}}$	M1		
$\theta - 17.354^\circ = 44.286^\circ$			
$\theta = 61.6^{\circ}$ (to 1 dec pl)	A1		

(ii) Find the value of θ for which OT = 1.2

[5]