


Name:		Index Number:		Class:	
-------	--	---------------	--	--------	--



**CATHOLIC HIGH SCHOOL**  
**Preliminary Examination**  
**Secondary 4 (O-Level Programme)**



**Additional Mathematics**

**4049/01**

Paper 1

**14 September 2021**

**2 hours 15 minutes**

Additional Materials: Answer Booklets A and B

**READ THESE INSTRUCTIONS FIRST**

Write your name, index number and class on all the work you hand in.

Write in dark blue or black pen.

You may use a soft pencil for any diagrams or graphs.

Do not use staples, paper clips, highlighters, glue or correction fluid.

Answer **all** questions in the space provided

If working is needed for any question, it must be shown with the answer.

Omission of essential working will result in loss of marks.

The use of an approved scientific calculator is expected, where appropriate.

If the degree of accuracy is not specified in the question, and if the answer is not exact, give the answer to **three significant figures**.

Give answers in **degrees to one decimal place**.

The number of marks is given in brackets [ ] at the end of each question or part question.

The total of the marks for this paper is **90**.

*For examiner's use:*

Q1	Q2	Q3	Q4	Q5	Q6	Total Marks
/ 9	/ 11	/ 9	/ 8	/ 8	/ 5	/ 50

This booklet consists of **12** printed pages.

**Mathematical Formulae****1. ALGEBRA***Quadratic Equation*

For the quadratic equation  $ax^2 + bx + c = 0$ ,

$$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$

*Binomial Expansion*

$$(a+b)^n = a^n + \binom{n}{1}a^{n-1}b + \binom{n}{2}a^{n-2}b^2 + \dots + \binom{n}{r}a^{n-r}b^r + \dots + b^n,$$

where  $n$  is a positive integer and  $\binom{n}{r} = \frac{n!}{(n-r)!r!} = \frac{n(n-1)\dots(n-r+1)}{r!}$

**2. TRIGONOMETRY***Identities*

$$\sin^2 A + \cos^2 A = 1$$

$$\sec^2 A = 1 + \tan^2 A$$

$$\operatorname{cosec}^2 A = 1 + \cot^2 A$$

$$\sin(A \pm B) = \sin A \cos B \pm \cos A \sin B$$

$$\cos(A \pm B) = \cos A \cos B \mp \sin A \sin B$$

$$\tan(A \pm B) = \frac{\tan A \pm \tan B}{1 \mp \tan A \tan B}$$

$$\sin 2A = 2 \sin A \cos A$$

$$\cos 2A = \cos^2 A - \sin^2 A = 2\cos^2 A - 1 = 1 - 2\sin^2 A$$

$$\tan 2A = \frac{2 \tan A}{1 - \tan^2 A}$$

*Formulae for  $\Delta ABC$* 

$$\frac{a}{\sin A} = \frac{b}{\sin B} = \frac{c}{\sin C}$$

$$a^2 = b^2 + c^2 - 2bc \cos A$$

$$\Delta = \frac{1}{2} ab \sin C$$

1. (a) Without using tables or calculator, find the value of  $k$  such that

$$\left(\frac{10}{\sqrt{15}} - \frac{\sqrt{243}}{3\sqrt{5}} + \frac{20}{\sqrt{180}}\right) \times \frac{3}{\sqrt{5}} = 2 + k\sqrt{3}.$$

[3]

- (b) Given that  $\log_4 xy = 7$  and  $\frac{\log_4 x}{\log_4 y} = -8$ , find the value of  $\log_4 y$ .

Hence, evaluate  $\log_4 \frac{2x}{y^3}$ .

[6]



2. The population of polar bears in the arctic is given by the formula

$$N = 8000(2 + 3e^{-\frac{t}{50}}), \text{ where } t \text{ is measured in years. Find}$$

(i) the initial population, [1]


(ii) the population after 50 years, [1]

(iii) the least number of years it would take the population to exceed 20 000, [3]

- (iv) the rate at which the polar bears is decreasing when  $t = 10$ , [2]

- (v) From the formula  $N = 8000(2 + 3e^{-\frac{t}{50}})$ , explain why the population of the polar bears can never fall below 16000. [2]

- (vi) Sketch the population-time curve in the grid below. [2]



3. Given that the expression  $2x^3 + ax^2 + bx - 6$  is divisible by  $(x^2 - 2)$ , find the value of  $a$  and of  $b$ . [4]

Hence,

- (i) solve the equation  $2x^3 + ax^2 + bx - 6 = 0$ , for the exact value(s) of  $x$ , [3]

- (ii) solve the equation  $2 + ay + by^2 - 6y^3 = 0$ . [2]

4. (a) Solve  $2\sin x = \tan x$  for which  $0^\circ \leq x \leq 360^\circ$ . [4]

- (b) Find all the angles between 0 and 6 for which  $2\tan y(\tan y + 1) = 3(2 - \sec^2 y)$ . [4]

5. (a) Given that  $\cos A = -\frac{3}{5}$  where  $\pi < A < \frac{3\pi}{2}$ . Find, without using a calculator, the value of

(i)  $\sin(\pi - A)$ , [1]

(ii)  $\cot A$ , [1]

(iii)  $\cos \frac{A}{2}$ . [2]

(b) Prove the identity

$$\frac{2 - \sec^2 x}{\sec^2 x + 2 \tan x} \equiv \frac{\cos x - \sin x}{\cos x + \sin x}$$

[4]

- 6 The function  $h$  is defined, for  $0^\circ \leq x \leq 360^\circ$ , by  $h(x) = a \cos(bx) + c$ , where  $a$ ,  $b$  and  $c$  are positive integers. Given that the greatest and least values of  $h$  are 3 and  $-1$  respectively and the period of  $h$  is  $720^\circ$ ,

(i) state the value of  $a$ , of  $b$  and of  $c$ . [3]

(ii) Sketch the graph of  $h$ . [2]



Name:		Index Number:		Class:	
-------	--	---------------	--	--------	--



**CATHOLIC HIGH SCHOOL**  
**Preliminary Examination**  
**Secondary 4 (O-Level Programme)**

**B**

**ADDITIONAL MATHEMATICS**

**4049/01**

Paper 1  
 BOOKLET B

**14 September 2021**  
**2 hours 15 minutes**

*For examiner's use:*

Q7	Q8	Q9	Q10	Q11	Total Marks
/ 6	/ 9	/ 9	/ 6	/ 10	/ 40

This Booklet B consists of **10** printed pages.

- 7 Express  $\frac{4x^3 + 7x^2 + 49x + 14}{(2x+1)(x^2+9)}$  in partial fractions. [6]

- 8 Two points  $A$  and  $B$  have coordinates  $(1, 2)$  and  $(-3, 6)$  respectively. A point  $P(x, y)$  is such that  $AP$  and  $BP$  are perpendicular.
- (a) Show that  $P$  lies on the circumference of a circle. [2]

Find

- (b) the coordinates of the centre of the circle and the radius of the circle, [4]


- (c) the equation of the circle. [1]

- (d) Explain why the tangents to the circle at  $A$  and  $B$  are parallel. [2]

- 9 (a) A curve has the equation  $y = \frac{2x+11}{x+1}$ ,  $x \neq -1$ . Show that  $y$  is a decreasing function. [3]

- (b) Find the coordinates of the stationary point(s) of the curve  $y = xe^{-2x}$  and determine the nature of the stationary point(s). [6]

- 10 The diagram shows part of the curve  $y = 4 \cos \frac{1}{3}x$ . Find the area of the shaded region bounded by the curve, the  $y$ -axis and the line  $y = -2$ . [6]






- 11 A particle  $A$  moves in a straight line, so that,  $t$  seconds after leaving a fixed point  $P$ , its velocity,  $v_A \text{ ms}^{-1}$ , is given by  $v_A = 2t - 6$ .

(a) Find the distance travelled by the particle  $A$  before it comes to instantaneous rest. [3]



A second particle  $B$  moves along the same horizontal line as  $A$ , and starts from  $Q$ , a point 8 m away from  $P$ , at the same instant that  $A$  begins to move. Particle  $B$  moves with a velocity of  $5 \text{ ms}^{-1}$  and decelerates at  $1 \text{ ms}^{-2}$ .



- (b) Calculate the distance between particle  $A$  and particle  $B$  when  $A$  comes to instantaneous rest. [3]

- (c) Find the time during the interval  $0 \leq t \leq 5$  when the distance between particle  $A$  and particle  $B$  is at its maximum. Calculate this maximum distance. [3]

- (d) Find the range of values of  $t$  for which both particles are moving in the same direction. [1]

**END OF PAPER**

ANS

Name:		Index Number:		Class:	
-------	--	---------------	--	--------	--

**MARKING SCHEME**



**CATHOLIC HIGH SCHOOL**  
**Preliminary Examination**  
**Secondary 4 (O-Level Programme)**

**A**

**Additional Mathematics**

**4049/01**

Paper 1

**14 September 2021**

**2 hours 15 minutes**

Additional Materials: Answer Booklets A and B

**READ THESE INSTRUCTIONS FIRST**

Write your name, index number and class on all the work you hand in.

Write in dark blue or black pen.

You may use a soft pencil for any diagrams or graphs.

Do not use staples, paper clips, highlighters, glue or correction fluid.

Answer **all** questions in the space provided

If working is needed for any question, it must be shown with the answer.

Omission of essential working will result in loss of marks.

The use of an approved scientific calculator is expected, where appropriate.

If the degree of accuracy is not specified in the question, and if the answer is not exact, give the answer answer to **three significant figures**.

Give answers in **degrees to one decimal place**.

For  $\pi$ , use either your calculator value or 3.142, unless the question requires the answer in terms of  $\pi$ .

The number of marks is given in brackets [ ] at the end of each question or part question.

The total of the marks for this paper is **90**.

This document consists of **18** printed pages, including 1 blank page.

## **Mathematical Formulae**

### **1. ALGEBRA**

#### *Quadratic Equation*

For the quadratic equation  $ax^2 + bx + c = 0$ ,

$$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$

#### *Binomial Expansion*

$$(a + b)^n = a^n + \binom{n}{1}a^{n-1}b + \binom{n}{2}a^{n-2}b^2 + \dots + \binom{n}{r}a^{n-r}b^r + \dots + b^n,$$

where  $n$  is a positive integer and  $\binom{n}{r} = \frac{n!}{(n-r)!r!} = \frac{n(n-1)\dots(n-r+1)}{r!}$

### **2. TRIGONOMETRY**

#### *Identities*

$$\sin^2 A + \cos^2 A = 1$$

$$\sec^2 A = 1 + \tan^2 A$$

$$\operatorname{cosec}^2 A = 1 + \cot^2 A$$

$$\sin(A \pm B) = \sin A \cos B \pm \cos A \sin B$$

$$\cos(A \pm B) = \cos A \cos B \mp \sin A \sin B$$

$$\tan(A \pm B) = \frac{\tan A \pm \tan B}{1 \mp \tan A \tan B}$$

$$\sin 2A = 2 \sin A \cos A$$

$$\cos 2A = \cos^2 A - \sin^2 A = 2\cos^2 A - 1 = 1 - 2\sin^2 A$$

$$\tan 2A = \frac{2 \tan A}{1 - \tan^2 A}$$

#### *Formulae for $\Delta ABC$*

$$\frac{a}{\sin A} = \frac{b}{\sin B} = \frac{c}{\sin C}$$

$$a^2 = b^2 + c^2 - 2bc \cos A$$

$$\Delta = \frac{1}{2}ab \sin C$$

1. (a) Without using tables or calculator, find the value of  $k$  such that

$$\left(\frac{10}{\sqrt{15}} - \frac{\sqrt{243}}{3\sqrt{5}} + \frac{20}{\sqrt{180}}\right) \times \frac{3}{\sqrt{5}} = 2 + k\sqrt{3}. \quad [3]$$

1(a)	$\begin{aligned} & \left(\frac{10}{\sqrt{15}} - \frac{\sqrt{243}}{3\sqrt{5}} + \frac{20}{\sqrt{180}}\right) \times \frac{3}{\sqrt{5}} \\ &= \frac{30}{5\sqrt{3}} - \frac{27\sqrt{3}}{15} + \frac{60}{30} \\ &= \frac{6}{\sqrt{3}} - \frac{9\sqrt{3}}{5} + 2 \\ &= \frac{30 - 27 + 10\sqrt{3}}{5\sqrt{3}} \\ &= \frac{3 + 10\sqrt{3}}{5\sqrt{3}} \times \frac{\sqrt{3}}{\sqrt{3}} \\ &= \frac{3\sqrt{3} + 30}{15} \\ &= 2 + \frac{1}{5}\sqrt{3} \\ &k = \frac{1}{5} \end{aligned}$
------	---

- (b) Given that  $\log_4 xy = 7$  and  $\frac{\log_4 x}{\log_4 y} = -8$ , find the value of  $\log_4 y$ .

Hence, evaluate  $\log_4 \frac{2x}{y^3}$ .

[6]

(b)	$\log_4 xy = 7$ $\log_4 x + \log_4 y = 7 \dots (1)$ $\log_4 x = -8 \log_4 y \dots (2)$ <p>Sub (2) into (1):</p> $-8 \log_4 y + \log_4 y = 7$ $\log_4 y = -1$ <p>Hence from (2),</p> $\log_4 x = 8$ $\log_4 \frac{2x}{y^3} = \log_4 2 + \log_4 x - 3 \log_4 y$ $= \frac{1}{\log_2 4} + 8 - 3(-1)$ $= 11 \frac{1}{2}$
-----	--

2. The population of polar bears in the arctic is given by the formula

$$N = 8000(2 + 3e^{-\frac{t}{50}}), \text{ where } t \text{ is measured in years. Find}$$

- (i) the initial population,

[1]

(i)	When $t = 0$ , $N = 8000(2 + 3e^0)$ $= 40\,000$
-----	--

- (ii) the population after 50 years,

[1]

(ii)	When $t = 50$ , $N = 8000(2 + 3e^{-\frac{t}{50}})$ $= 8000(2 + 3e^{-\frac{50}{50}})$ $= 8000(2 + 3e^{-1})$ $= 24829.10$ $= 24800$
------	--

- (iii) the least number of years it would take the population to exceed 20 000,

[3]

(iii)	$20000 = 8000(2 + 3e^{-\frac{t}{50}})$ $(2 + 3e^{-\frac{t}{50}}) = \frac{20000}{8000}$ $e^{-\frac{t}{50}} = \frac{1}{6}$ $-\frac{t}{50} = \ln \frac{1}{6}$ $t = -50 \ln \frac{1}{6}$ $= 89.587$ $= 89.6$ <p>Hence it will take at least 90 years.</p>
-------	---

- (iv) the rate at which the polar bears is decreasing when  $t = 10$ , [2]

(iv)	$N = 8000(2 + 3e^{-\frac{t}{50}})$ $\frac{dN}{dt} = 24000\left(-\frac{1}{50}\right)e^{-\frac{t}{50}}$ $= -480e^{-\frac{t}{50}}$ <p>When <math>t = 10</math></p> $\frac{dN}{dt} = -480e^{-\frac{10}{50}}$ $= -392.99$ $= -393$ <p>The population of the polar bears is decreasing at a rate of 393 polar bears /year.</p>
------	--


- (v) From the formula  $N = 8000(2 + 3e^{-\frac{t}{50}})$ , explain why the population of the polar bears can never fall below 16000. [2]

(v)	<p>As <math>t \rightarrow \infty</math>,</p> $e^{-\frac{t}{50}} \rightarrow 0$ $\therefore N \rightarrow 8000(2) = 16000$ <p>Hence the population of the polar bears will never fall below 16000.</p> <p><b>Alternative Method</b> Use graph to explain</p>
-----	---



- (vi) Sketch the population-time curve in the grid below.

[2]



3. Given that the expression  $2x^3 + ax^2 + bx - 6$  is divisible by  $(x^2 - 2)$ , find the value of  $a$  and of  $b$ .

[4]

$$\text{Let } f(x) = 2x^3 + ax^2 + bx - 6$$

$$f(\sqrt{2}) = 2(\sqrt{2})^3 + a(\sqrt{2})^2 + b(\sqrt{2}) - 6 = 0$$

$$4(\sqrt{2}) + 2a + (\sqrt{2})b - 6 = 0 \dots (1)$$

$$f(-\sqrt{2}) = -2(\sqrt{2})^3 + a(\sqrt{2})^2 - b(\sqrt{2}) - 6 = 0$$

$$-4(\sqrt{2}) + 2a - (\sqrt{2})b - 6 = 0 \dots (2)$$

$$(1) + (2)$$

$$4a - 12 = 0$$

$$a = 3$$

$$(1) - (2)$$

$$8(\sqrt{2}) + 2(\sqrt{2})b = 0$$

$$b = \frac{-8(\sqrt{2})}{2(\sqrt{2})} = -4$$

**Alternative Method**

$$\text{Let } f(x) = 2x^3 + ax^2 + bx - 6 = (x^2 - 2)(px + q)$$

$$2x^3 + ax^2 + bx - 6 = px^3 + qx^2 - 2px - 2q$$

$$\left\{ \begin{array}{l} \text{Comparing coefficients of } x^3 : p = 2 \\ \text{Comparing constant terms: } q = 3 \end{array} \right\}$$

$$\therefore a = 3$$

$$b = -2(2)$$

$$= -4$$

Hence,

- (i) solve the equation  $2x^3 + ax^2 + bx - 6 = 0$ , giving the exact value(s) of  $x$ . [3]

(i)  $f(x) = 2x^3 + 3x^2 - 4x - 6 = 0$

$$(x^2 - 2)(2x + 3) = 0$$

$$(x + \sqrt{2})(x - \sqrt{2})(2x + 3) = 0$$

$$x = -\sqrt{2} \text{ or } x = \sqrt{2} \text{ or } x = -\frac{3}{2}$$

Alternative Method

$$2x^3 + 3x^2 - 4x - 6 = 0$$

$$(x^2 - 2)(px + q) = 0$$

$$\text{Comparing coefficients of } x^3 : p = 2$$

$$\text{Comparing constant terms : } -2q = -6$$

$$q = 3$$

$$f(x) = 2x^3 + 3x^2 - 4x - 6 = 0$$

$$(x^2 - 2)(2x + 3) = 0$$

$$(x + \sqrt{2})(x - \sqrt{2})(2x + 3) = 0$$

$$x = -\sqrt{2} \text{ or } x = \sqrt{2} \text{ or } x = -\frac{3}{2}$$

(ii) solve the equation  $2 + ay + by^2 - 6y^3 = 0$ .

[2]

(ii)	$2 + 3y - 4y^2 - 6y^3 = 0$ $2\left(\frac{1}{y}\right)^3 + 3\left(\frac{1}{y}\right)^2 - 4\left(\frac{1}{y}\right) - 6 = 0$ $\therefore \frac{1}{y} = x$ $\frac{1}{y} = -\sqrt{2}, \sqrt{2}, -\frac{3}{2}$ $y = -\frac{1}{\sqrt{2}} \text{ or } y = \frac{1}{\sqrt{2}} \text{ or } y = -\frac{2}{3}$
------	---

4. (a) Solve  $2 \sin x = \tan x$  for which  $0^\circ \leq x \leq 360^\circ$ . [4]

(a)	$2 \sin x = \tan x$ $2 \sin x = \frac{\sin x}{\cos x}$ $2 \sin x \cos x - \sin x = 0$ $\sin x(2 \cos x - 1) = 0$ $\sin x = 0 \quad \text{or} \quad \cos x = \frac{1}{2}$ $x = 0^\circ, 180^\circ, 360^\circ \quad \text{or} \quad x = 60^\circ, 300^\circ$ $x = 0^\circ, 60^\circ, 180^\circ, 300^\circ, 360^\circ$
-----	---

- (b) Find all the angles between 0 and 6 for which  $2 \tan y(\tan y + 1) = 3(2 - \sec^2 y)$ . [4]

(b)	$2 \tan y(\tan y + 1) = 3(2 - \sec^2 y)$ $2 \tan^2 y + 2 \tan y = 6 - 3(1 + \tan^2 y)$ $5 \tan^2 y + 2 \tan y - 3 = 0$ $(5 \tan y - 3)(\tan y + 1) = 0$ $\tan y = \frac{3}{5} \quad \text{or} \quad \tan y = -1$ $y = 0.540, \pi + 0.5404195$ $= 0.540, 3.68$ $y = 0.540, \frac{3\pi}{4}, 3.68, \frac{7\pi}{4}$ <p>or</p> $y = 0.540, 2.36, 3.68, 5.50$	$BA = \frac{\pi}{4} \text{ or } 0.785398$ $y = \pi - \frac{\pi}{4}, 2\pi - \frac{\pi}{4}$ $= \frac{3\pi}{4}, \frac{7\pi}{4} \text{ or } 2.36, 5.50$
-----	---	---

5. (a) Given that  $\cos A = -\frac{3}{5}$  where  $\pi < A < \frac{3\pi}{2}$ . Find, without using a calculator, the value of

(i)  $\sin(\pi - A)$ , [1]

(i)	$\sin(\pi - A) = \sin A$ $= -\frac{4}{5}$
-----	---

(ii)  $\cot A$ , [1]

(ii)	$\cot A = \frac{1}{\tan A}$ $= \frac{3}{4}$
------	---

(iii)  $\cos \frac{A}{2}$ . [2]

(iii)	$\cos A = 2\cos^2 \frac{A}{2} - 1$ $\cos \frac{A}{2} = -\sqrt{\frac{1 + \cos A}{2}}$ $= -\sqrt{\frac{1 - \frac{3}{5}}{2}}$ $= -\sqrt{\frac{1}{5}}$ $= -\frac{\sqrt{5}}{5}$
-------	--

(b) Prove the identity

$$\frac{2 - \sec^2 x}{\sec^2 x + 2 \tan x} \equiv \frac{\cos x - \sin x}{\cos x + \sin x}$$

[4]

(b)

$$\begin{aligned} LHS &= \frac{2 - \sec^2 x}{\sec^2 x + 2 \tan x} \\ &= \frac{2 - \frac{1}{\cos^2 x}}{\frac{1}{\cos^2 x} + \frac{2 \sin x}{\cos x}} \\ &= \frac{2 \cos^2 x - 1}{1 + 2 \sin x \cos x} \\ &= \frac{\cos^2 x - \sin^2 x}{\sin^2 x + \cos^2 x + 2 \sin x \cos x} \\ &= \frac{(\cos x + \sin x)(\cos x - \sin x)}{(\cos x + \sin x)^2} \\ &= \frac{\cos x - \sin x}{\cos x + \sin x} \equiv RHS \end{aligned}$$

Alternative Method

$$\begin{aligned} &\frac{2 - (1 + \tan^2 x)}{1 + \tan^2 x + 2 \tan x} \\ &= \frac{1 - \tan^2 x}{(1 + \tan x)^2} \\ &= \frac{(1 - \tan x)(1 + \tan x)}{(1 + \tan x)^2} \\ &= \frac{(1 - \tan x)}{(1 + \tan x)} \\ &= \frac{1 - \frac{\sin x}{\cos x}}{1 + \frac{\sin x}{\cos x}} \\ &= \frac{\frac{\cos x - \sin x}{\cos x}}{\frac{\cos x + \sin x}{\cos x}} \\ &= \frac{\cos x - \sin x}{\cos x + \sin x} \end{aligned}$$

- 6 The function  $h$  is defined, for  $0^\circ \leq x \leq 360^\circ$ , by  $h(x) = a \cos(bx) + c$ , where  $a$ ,  $b$  and  $c$  are positive integers. Given that the greatest and least values of  $h$  are 3 and  $-1$  respectively and the period of  $h$  is  $720^\circ$ ,


(i) state the value of  $a$ , of  $b$  and of  $c$ .

[3]

(i)	$a = 2$ $c = 1$  Period, $720^\circ = \frac{360^\circ}{b}$  $b = \frac{1}{2}$
-----	--

(ii) Sketch the graph of  $h$ .

[2]



- 7 Express  $\frac{4x^3 + 7x^2 + 49x + 14}{(2x+1)(x^2+9)}$  in partial fractions.

[6]

$$\begin{array}{r}
 2x^3 + x^2 + 18x + 9 \overline{) 4x^3 + 7x^2 + 49x + 14} \\
 \underline{4x^3 + 2x^2 + 36x + 18} \phantom{00} \\
 5x^2 + 13x - 4
 \end{array}$$

$$\frac{4x^3 + 7x^2 + 49x + 14}{(2x+1)(x^2+9)} = 2 + \frac{5x^2 + 13x - 4}{(2x+1)(x^2+9)}$$

$$\frac{5x^2 + 13x - 4}{(2x+1)(x^2+9)} = \frac{A}{2x+1} + \frac{Bx+C}{x^2+9}$$

$$\begin{aligned}
 5x^2 + 13x - 4 &= A(x^2+9) + (Bx+C)(2x+1) \\
 &= (A+2B)x^2 + (B+2C)x + 9A+C
 \end{aligned}$$

Comparing coefficients:

$$A + 2B = 5 \dots (1)$$

$$B + 2C = 13 \dots (2)$$

$$9A + C = -4 \dots (3)$$

$$A = -1$$

$$B = 3$$

$$C = 5$$

$$\frac{4x^3 + 7x^2 + 49x + 14}{(2x+1)(x^2+9)} = 2 - \frac{1}{2x+1} + \frac{3x+5}{x^2+9}$$



- 8 Two points  $A$  and  $B$  have coordinates  $(1, 2)$  and  $(-3, 6)$  respectively. A point  $P(x, y)$  is such that  $AP$  and  $BP$  are perpendicular.

(a) Show that  $P$  lies on the circumference of a circle.

[2]

(a)	<p>Since <math>AP \perp PB</math> (given), <math>\angle APB = 90^\circ</math>  Hence <math>P</math> lies on the circumference of a semi-circle  (Right angle in semi-circle)</p>
-----	--

Find

(b) the coordinates of the centre of the circle and the radius of the circle,

[4]

(b)	<p>From (a), we know <math>AB</math> is a diameter of the circle  Centre of circle,  <math>C\left(\frac{1-3}{2}, \frac{2+6}{2}\right)</math>  <math>= (-1, 4)</math>  Radius, <math>r = \sqrt{(1+1)^2 + (2-4)^2}</math>  <math>= \sqrt{8}</math>  <math>= 2\sqrt{2}</math></p>
-----	--

(c) the equation of the circle.

[1]

(c)	$(x+1)^2 + (y-4)^2 = (\sqrt{8})^2$ $(x+1)^2 + (y-4)^2 = 8$
-----	--

(d) Explain why the tangents to the circle at  $A$  and  $B$  are parallel.

[2]

(d)	<p>Since <math>AB</math> is a diameter of the circle,  both tangents to the circle at <math>A</math> and <math>B</math> are perpendicular to <math>AB</math>  (radius is perpendicular to the tangent)  Hence the tangents to the circle at <math>A</math> and <math>B</math> are parallel.</p>
-----	---


- 9 (a) A curve has the equation  $y = \frac{2x+11}{x+1}$ ,  $x \neq -1$ . Show that  $y$  is a decreasing function. [3]

(a)	$y = \frac{2x+11}{x+1}$ $\frac{dy}{dx} = \frac{(x+1)2 - (2x+11)}{(x+1)^2}$ $= \frac{-9}{(x+1)^2}$ <p>Since <math>(x+1)^2 &gt; 0</math> and <math>-9</math> is a negative constant,</p> $\frac{dy}{dx} < 0$ , hence $y$ is a decreasing function
-----	---

- (b) Find the coordinates of the stationary point(s) of the curve  $y = xe^{-2x}$  and determine the nature of the stationary point(s). [6]

(b)	$y = xe^{-2x}$ $\frac{dy}{dx} = -2xe^{-2x} + e^{-2x}$ $= e^{-2x}(1 - 2x)$ <p>For stationary points, <math>\frac{dy}{dx} = 0</math></p> $e^{-2x}(1 - 2x) = 0$ $x = \frac{1}{2}$ $y = \frac{1}{2e}$ <p><math>\therefore</math> the stationary point is <math>\left(\frac{1}{2}, \frac{1}{2e}\right)</math>.</p> $\frac{d^2y}{dx^2} = -2e^{-2x} - 2(1 - 2x)e^{-2x}$ $= 4e^{-2x}(x - 1)$ <p>When <math>x = \frac{1}{2}</math>,</p> $\frac{d^2y}{dx^2} = 4e^{-1}\left(\left(\frac{1}{2}\right) - 1\right) < 0$ <p><math>\left(\frac{1}{2}, \frac{1}{2e}\right)</math> is a maximum point.</p>
-----	--

- 10 The diagram shows part of the curve  $y = 4 \cos \frac{1}{3}x$ . Find the area of the shaded region bounded by the curve, the  $y$ -axis and the line  $y = -2$ . [6]



$$y = 4 \cos \frac{1}{3}x$$

$$4 \cos \frac{1}{3}x = -2 \quad \text{and} \quad 2 \cos \frac{1}{3}x = 0$$

$$\cos \frac{1}{3}x = -\frac{1}{2} \quad \frac{1}{3}x = \frac{\pi}{2}$$

$$BA = \frac{\pi}{3} \quad x = \frac{3\pi}{2}$$

$$\frac{1}{3}x = \frac{2\pi}{3}$$

$$x = 2\pi$$

$$\text{Area of the shaded region} = \int_0^{2\pi} \left( 4 \cos \frac{1}{3}x + 2 \right) dx$$

$$= \left[ 12 \sin \frac{1}{3}x + 2x \right]_0^{2\pi}$$

$$= \left( 12 \sin \frac{2\pi}{3} + 2(2\pi) - 0 \right)$$

$$= 12 \frac{\sqrt{3}}{2} + 4\pi$$

$$= (6\sqrt{3} + 4\pi) \text{ units}^2$$

$$= 22.95$$

$$= 23.0 \text{ units}^2$$

Alternative Method

Area of the shaded region

$$= \int_0^{\frac{3\pi}{2}} \left( 4 \cos \frac{1}{3}x \right) dx + 2(2\pi) - \left[ -\int_{\frac{3\pi}{2}}^{2\pi} \left( 4 \cos \frac{1}{3}x \right) dx \right]$$

$$= \left[ 12 \sin \frac{1}{3}x \right]_0^{\frac{3\pi}{2}} + 4\pi + \left[ 12 \sin \frac{1}{3}x \right]_{\frac{3\pi}{2}}^{2\pi}$$

$$= 12 \sin \frac{\pi}{2} - 0 + 4\pi + 12 \sin \frac{2\pi}{3} - 12 \sin \frac{\pi}{2}$$

$$= 4\pi + 12 \frac{\sqrt{3}}{2}$$

$$= (6\sqrt{3} + 4\pi) \text{ units}^2$$

$$= 23.0 \text{ units}^2$$


- 11 A particle  $A$  moves in a straight line, so that,  $t$  seconds after leaving a fixed point  $P$ , its velocity,  $v_A \text{ ms}^{-1}$ , is given by  $v_A = 2t - 6$ .

(a) Find the distance travelled by the particle  $A$  before it comes to instantaneous rest.

[3]

(a)	<p>At rest, <math>V_A = 2t - 6 = 0</math> [M1] <math>V_A = 0</math>  <math>\therefore t = 3</math></p> <div style="display: flex; justify-content: space-between;"> <div style="width: 45%;"> <math>s_A = \int (2t - 6) dt</math>  <math>= t^2 - 6t + c</math>  <math>t = 0, s = 0 \Rightarrow c = 0</math>  <math>\therefore s_A = t^2 - 6t</math>            When <math>t = 3, s_A = 3^2 - 6(3) = -9m</math> </div> <div style="width: 45%;"> <math>\text{or } s_A = \int_0^3 (2t - 6) dt</math>  <math>= \left[ t^2 - 6t \right]_0^3</math>  <math>= 3^2 - 6(3) = -9</math> </div> </div> <p style="text-align: center;">Hence total distance travelled is 9 m</p>
-----	---

A second particle  $B$  moves along the same horizontal line as  $A$ , and at the same instant that  $A$  begins to move.  $B$  starts from  $Q$ , a point 8 m away from  $P$ , with a velocity of  $5 \text{ ms}^{-1}$  and decelerates at  $1 \text{ ms}^{-2}$ .



- (b) Calculate the distance between particle  $A$  and particle  $B$  when  $A$  comes to instantaneous rest. [3]

(b)	<div style="display: flex; justify-content: space-between;"> <div style="width: 45%;"> <math display="block">s_B = \int (5 - t) dt</math> <math display="block">= 5t - \frac{t^2}{2} + c</math> <math display="block">t = 0, s = 8 \Rightarrow c = 8</math> <math display="block">\therefore s_B = 5t - \frac{t^2}{2} + 8</math> <math display="block">\text{When } t = 3, s_B = 15 - \frac{9}{2} + 8 = 18\frac{1}{2} \text{ m}</math> <math display="block">\text{Distance between A and B} = 18\frac{1}{2} + 9</math> <math display="block">= 27.5 \text{ m}</math> </div> <div style="width: 45%;"> <math display="block">\text{or } s_B = \int_0^3 (5 - t) dt</math> <math display="block">= \left[ 5t - \frac{t^2}{2} \right]_0^3</math> <math display="block">= 5(3) - \frac{9}{2}</math> <math display="block">= 10\frac{1}{2}</math> <math display="block">\text{Distance between A and B} = 10\frac{1}{2} + 8 + 9</math> <math display="block">= 27.5 \text{ m}</math> </div> </div>
-----	---

- (c) Find the time during the interval  $0 \leq t \leq 5$  when the distance between particle  $A$  and particle  $B$  is at its maximum. Calculate this maximum distance. [3]

(c)	$s_B = 5t - \frac{t^2}{2} + 8$ $s_{AB} = 11t - \frac{3t^2}{2} + 8$ $\frac{ds}{dt} = 11 - 3t$ <p>For max distance, <math>\frac{ds}{dt} = 0</math></p> $\therefore t = \frac{11}{3}$ <p>When <math>t = \frac{11}{3}</math>,</p> $s = 11\left(\frac{11}{3}\right) - \frac{3}{2}\left(\frac{11}{3}\right)^2 + 8$ <p>Max distance is <math>= 28\frac{1}{6} m</math></p>
-----	--

- (d) Find the range of values of  $t$  for which both particles are moving in the same direction. [1]

(d)	<p>Hence both particles are moving in the same direction for the period <math>3 &lt; t &lt; 5</math>.</p>
-----	---

END OF PAPER



Name:		Index Number:		Class:	
-------	--	---------------	--	--------	--



**CATHOLIC HIGH SCHOOL**  
**2021 Preliminary Examination**  
**Secondary 4 (O-Level Programme)**

**ADDITIONAL MATHEMATICS**

**4049/02**

Paper 2

**15 September 2021**

**2 hours 15 minutes**

Additional materials: Answer booklets A, B and C

**READ THESE INSTRUCTIONS FIRST**

Write your name, register number and class on all the work you hand in.

Write in dark blue or black pen.

You may use a soft pencil for any diagrams or graphs.

Do not use staples, paper clips, highlighters, glue or correction fluid.

**DO NOT WRITE ON THE MARGINS.**

Answer **all** questions.

If working is needed for any question, it must be shown with the answer.

Omission of essential working will result in loss of marks.

The use of an approved scientific calculator is expected, where appropriate.

If the degree of accuracy is not specified in the question, and if the answer is not exact, give the answer to **three significant figures**. Give answers in **degrees to one decimal place**.

For  $\pi$ , use either your calculator value or 3.142, unless the question requires the answer in terms of  $\pi$ .

At the end of the examination, fasten all your work securely together.

The number of marks is given in brackets [ ] at the end of each question or part question.

The total of the marks for this paper is **90**.

This document consists of **22** printed pages, including 1 blank page.

[TURN OVER

**Mathematical Formulae****1. ALGEBRA***Quadratic Equation*

For the quadratic equation  $ax^2 + bx + c = 0$ ,

$$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$

*Binomial Expansion*

$$(a + b)^n = a^n + \binom{n}{1}a^{n-1}b + \binom{n}{2}a^{n-2}b^2 + \dots + \binom{n}{r}a^{n-r}b^r + \dots + b^n,$$

where  $n$  is a positive integer and  $\binom{n}{r} = \frac{n!}{(n-r)!r!} = \frac{n(n-1)\dots(n-r+1)}{r!}$

**2. TRIGONOMETRY***Identities*

$$\sin^2 A + \cos^2 A = 1$$

$$\sec^2 A = 1 + \tan^2 A$$

$$\operatorname{cosec}^2 A = 1 + \cot^2 A$$

$$\sin(A \pm B) = \sin A \cos B \pm \cos A \sin B$$

$$\cos(A \pm B) = \cos A \cos B \mp \sin A \sin B$$

$$\tan(A \pm B) = \frac{\tan A \pm \tan B}{1 \mp \tan A \tan B}$$

$$\sin 2A = 2 \sin A \cos A$$

$$\cos 2A = \cos^2 A - \sin^2 A = 2\cos^2 A - 1 = 1 - 2\sin^2 A$$

$$\tan 2A = \frac{2 \tan A}{1 - \tan^2 A}$$

*Formulae for  $\Delta ABC$* 

$$\frac{a}{\sin A} = \frac{b}{\sin B} = \frac{c}{\sin C}$$

$$a^2 = b^2 + c^2 - 2bc \cos A$$

$$\Delta = \frac{1}{2}ab \sin C$$

- 1 (a) Solve the simultaneous equations

$$2y - 7x = 5x^2,$$

$$3y = 7 - 4x.$$


[3]

- (b) Given that the line  $y = mx + 1$  is a tangent to the curve  $y = x^2 + m(2x + 1)$ , find the value(s) of the constant  $m$ .

[3]

- (c) Find the range of values of  $k$  such that  $x^2 + 4x > kx - 9$ . [3]

- 2 The diagram shows a quadrilateral with vertices  $O(0, 0)$ ,  $A(8, -2)$ ,  $B$  and  $C$ . The equation of  $AB$  is  $y = 4x - 34$  and the equation of  $BC$  is  $5y + 3x = 60$ .



- (i) Find the coordinates of  $B$  and  $C$ . [3]

(ii) Determine if  $OABC$  is a kite.


[2]

(iii) A point  $D$ , above the  $x$ -axis, is such that  $OD$  is parallel to  $AB$  and  $OD = 5\sqrt{17}$ , find the coordinates of  $D$ .

[4]

- (iv) Find the area of the trapezium  $OABD$ . [2]

- 3 The diagram shows a structure such that  $VA = 3$  m,  $AB = 2$  m,  $BC = 1.5$  m and angle  $AVD = \text{angle } ABF = \theta$  degrees.  $BF$  is parallel to  $CE$ ,  $D$  is the midpoint of  $CE$  and the structure is symmetrical about  $VD$ .



- (i) Show that  $VD = 3 \cos \theta + 2 \sin \theta + 1.5$ .

[1]

- (ii) Express  $VD$  in the form  $R \cos(\theta - \alpha) + 1.5$ , where  $R > 0$  and  $\alpha$  is acute.

[3]

(iii) State the largest value of  $VD$  and find the corresponding value of  $\theta$ . [2]

(iv) Find the value(s) of  $\theta$  for which  $VD = 5$  m. [4]



4 (a) Solve the equation  $8e^{x+1} = \frac{15}{e^{x-1}} - 14e$ .

[4]

(b) Solve the equation  $3\sin 2\theta = \cos \theta$  for  $-\pi \leq \theta \leq \pi$ .

[4]

5 (a) Solve the equation  $\log_5 x + 1 = 2\log_x 5$ .

[4]

(b) Solve the equation  $\log_9(2+x) = \log_{16} 4 - \log_{\frac{1}{3}} \sqrt{1-2x}$ . [5]

- 6 (i) Show that  $\frac{d}{dx} [x\sqrt{3x-1}] = \frac{Ax+B}{\sqrt{3x-1}}$ , where  $A$  and  $B$  are constants. [4]

- (ii) Hence or otherwise, find  $\int_1^2 \frac{9x-2}{\sqrt{3x-1}} dx$ , giving your answer correct to 3 significant figures. [4]


- 7 The function  $f$  is defined for  $x \neq -\frac{1}{3}$  and is such that  $f''(x) = 4e^{2x} + \frac{9}{(3x+1)^2}$ .

Given that  $f'(0) = -1$  and  $f(0) = 2$ , find an expression for  $f(x)$ . [7]

- 8 (a) The equation of a curve is  $y = \frac{1}{\sqrt{10x-1}} + \frac{1}{3}x$ , for  $x \neq \frac{1}{10}$ . Find the equation of the normal to the curve at the point  $\left(1, \frac{2}{3}\right)$ . [4]



- (b) The diagram shows a cylindrical water tank with base radius 16 cm.



Water is flowing into the tank at a constant rate of  $4.8\pi k \text{ cm}^3/\text{min}$ , where  $k$  is a constant. At time  $t$  minutes, the depth of the water in the tank is  $h$  metres. There is a hole at point  $P$  at the bottom of the tank and a cork is used to stop water leakage. When the cork is unplugged, water will leak from the hole at a rate of  $6.4\pi \text{ cm}^3/\text{min}$ .

Find an expression for the rate of change of  $h$  in terms of  $k$  after the cork is unplugged. [4]

- 9 (i) The expansion of  $(1+px)^n$ , where  $n > 0$ , by the binomial theorem is  $1+36x+66p^2x^2+kx^3+\dots$ . Find the value of  $n$ , of  $p$  and of  $k$ . [5]

- (ii) Find the term independent of  $x$  in the expansion of  $(3+x)\left(2x-\frac{1}{x}\right)^8$ . [4]

- 10** The table shows, to 2 decimal places, the population,  $P$ , of a certain kind of bacteria in a test-tube, measured at hourly intervals,  $t$ .

$t$	0.2	0.6	0.9	1.3	1.5
$P$	0.72	0.96	1.40	1.55	1.78

- (i) On the grid on the next page, plot  $\lg P$  against  $t$  and draw a straight line graph. [3]

- (ii) Determine which value of  $P$ , in the table above, is the incorrect reading.  
Use the graph to estimate a value of  $P$  to replace the incorrect recording of  $P$ . [2]

- (iii) Find the gradient of your straight line and hence express  $P$  in the form  $\frac{A^t}{10^{2k}}$ ,  
where  $A$  and  $k$  are constants. [4]

- (iv) On the same diagram, draw the line representing  $P^{10} = 10^{3-t}$  and hence estimate the  
value of  $t$  for which  $\left(\frac{A^t}{10^{2k}}\right)^{10} = 10^{3-t}$ . [2]

**BLANK PAGE**

ANS

Name:		Index Number:		Class:	
-------	--	---------------	--	--------	--



**CATHOLIC HIGH SCHOOL**  
**2021 Preliminary Examination**  
**Secondary 4 (O-Level Programme)**

Full solutions

**ADDITIONAL MATHEMATICS**

**4049/02**

Paper 2

**15 September 2021**  
**2 hours 15 minutes**

Additional materials: Answer booklets A, B and C

**READ THESE INSTRUCTIONS FIRST**

Write your name, register number and class on all the work you hand in.

Write in dark blue or black pen.

You may use a soft pencil for any diagrams or graphs.

Do not use staples, paper clips, highlighters, glue or correction fluid.

**DO NOT WRITE ON THE MARGINS.**

Answer **all** questions.

If working is needed for any question, it must be shown with the answer.

Omission of essential working will result in loss of marks.

The use of an approved scientific calculator is expected, where appropriate.

If the degree of accuracy is not specified in the question, and if the answer is not exact, give the answer to **three significant figures**. Give answers in **degrees to one decimal place**.

For  $\pi$ , use either your calculator value or 3.142, unless the question requires the answer in terms of  $\pi$ .

At the end of the examination, fasten all your work securely together.

The number of marks is given in brackets [ ] at the end of each question or part question.

The total of the marks for this paper is **90**.

This document consists of **8** printed pages, including 1 blank page.

*Mathematical Formulae*

[TURN OVER

## 1. ALGEBRA

### Quadratic Equation

For the quadratic equation  $ax^2 + bx + c = 0$ ,

$$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$

### Binomial Expansion

$$(a+b)^n = a^n + \binom{n}{1}a^{n-1}b + \binom{n}{2}a^{n-2}b^2 + \dots + \binom{n}{r}a^{n-r}b^r + \dots + b^n,$$

where  $n$  is a positive integer and  $\binom{n}{r} = \frac{n!}{(n-r)!r!} = \frac{n(n-1)\dots(n-r+1)}{r!}$

## 2. TRIGONOMETRY

### Identities

$$\sin^2 A + \cos^2 A = 1$$

$$\sec^2 A = 1 + \tan^2 A$$

$$\operatorname{cosec}^2 A = 1 + \cot^2 A$$

$$\sin(A \pm B) = \sin A \cos B \pm \cos A \sin B$$

$$\cos(A \pm B) = \cos A \cos B \mp \sin A \sin B$$

$$\tan(A \pm B) = \frac{\tan A \pm \tan B}{1 \mp \tan A \tan B}$$

$$\sin 2A = 2 \sin A \cos A$$

$$\cos 2A = \cos^2 A - \sin^2 A = 2\cos^2 A - 1 = 1 - 2\sin^2 A$$

$$\tan 2A = \frac{2 \tan A}{1 - \tan^2 A}$$

### Formulae for $\Delta ABC$

$$\frac{a}{\sin A} = \frac{b}{\sin B} = \frac{c}{\sin C}$$

$$a^2 = b^2 + c^2 - 2bc \cos A$$

$$\Delta = \frac{1}{2}ab \sin C$$

- 1 (a) Solve the simultaneous equations

$$2y - 7x = 5x^2,$$

$$3y = 7 - 4x.$$

[3]

$$2y - 7x = 5x^2 \Rightarrow y = \frac{5}{2}x^2 + \frac{7}{2}x \dots (1)$$

$$3y = 7 - 4x \dots (2)$$

Sub (1) into (2):

$$3\left(\frac{5}{2}x^2 + \frac{7}{2}x\right) = 7 - 4x$$

$$15x^2 + 29x - 14 = 0$$

$$(5x - 2)(3x + 7) = 0$$

$$x = \frac{2}{5} \quad \text{or} \quad -\frac{7}{3}$$

$$y = \frac{9}{5} \quad \text{or} \quad \frac{49}{9}$$

- (b) Given that the line  $y = mx + 1$  is a tangent to the curve  $y = x^2 + m(2x + 1)$ , find the value(s) of the constant  $m$ .

[3]

Sub  $y = mx + 1$  into  $y = x^2 + m(2x + 1)$ :

$$x^2 + m(2x + 1) = mx + 1$$

$$x^2 + mx + m - 1 = 0$$

$$D = 0$$

$$m^2 - 4(1)(m - 1) = 0$$

$$m^2 - 4m + 4 = 0$$

$$(m - 2)^2 = 0$$

$$m = 2$$

- (c) Find the range of values of  $k$  such that  $x^2 + 4x > kx - 9$ .

[3]

$$x^2 + 4x > kx - 9$$

$$x^2 + (4 - k)x + 9 > 0$$

$$D < 0$$


$$(4 - k)^2 - 4(1)(9) < 0$$

$$k^2 - 8k - 20 < 0$$

$$-2 < k < 10$$



- 2 The diagram shows a quadrilateral with vertices  $O(0, 0)$ ,  $A(8, -2)$ ,  $B$  and  $C$ . The equation of  $AB$  is  $y = 4x - 34$  and the equation of  $BC$  is  $5y + 3x = 60$ .



- (i) Find the coordinates of  $B$  and  $C$ .

[3]

Sub  $x = 0$  into  $5y + 3x = 60$ :

$$y = 12$$

$$C(0, 12)$$

Sub  $y = 4x - 34$  into  $5y + 3x = 60$ :

$$5(4x - 34) + 3x = 60$$

$$23x = 230$$

$$x = 10$$

$$y = 6$$

$$B(10, 6)$$

- (ii) Determine if  $OABC$  is a kite.

[2]

**Mtd 1:**

$$\text{Grad}_{OB} = \frac{6}{10} = \frac{3}{5}$$

$$\text{Grad}_{AC} = \frac{-2 - 12}{8 - 0} = -\frac{7}{4}$$

$$\text{Grad}_{OB} \times \text{Grad}_{AC} = \frac{3}{5} \times \left(-\frac{7}{4}\right)$$

$$\neq -1$$

$\Rightarrow OB$  is not perpendicular to  $AC$ .

Hence,  $OABC$  is **NOT** a kite.

**Mtd 2:**

$$OC = 12$$

$$BC = \sqrt{(10-0)^2 + (6-12)^2}$$

$$= \sqrt{136} \text{ or } 11.7$$

$$BC \neq OC$$

Hence,  $OABC$  is **NOT** a kite.

- (iii) A point  $D$ , above the  $x$ -axis, is such that  $OD$  is parallel to  $AB$  and  $OD = 5\sqrt{17}$ , find the coordinates of  $D$ . [4]

Since  $AB$  is  $y = 4x - 34$

$$\text{Grad}_{AB} = 4$$

Eqn. of  $OD$ :  $y = 4x$

Let  $D(x, 4x)$ :

$$OD = 5\sqrt{17}$$

$$\sqrt{x^2 + (4x)^2} = 5\sqrt{17}$$

$$17x^2 = 25(17)$$

$$x^2 = 25$$

$$x = 5 \quad (x > 0 \text{ since } D \text{ is above } x\text{-axis, } y > 0)$$


$$D(5, 20)$$

- (iv) Find the area of the trapezium  $OABD$ . [2]

$$\text{area of the trapezium } OABD = \frac{1}{2} \begin{vmatrix} 0 & 8 & 10 & 5 & 0 \\ 0 & -2 & 6 & 20 & 0 \end{vmatrix}$$


$$= 119 \text{ units}^2$$

- 3 The diagram shows a structure such that  $VA = 3$  m,  $AB = 2$  m,  $BC = 1.5$  m and angle  $AVD = \text{angle } ABF = \theta$  degrees.  $BF$  is parallel to  $CE$ ,  $D$  is the midpoint of  $CE$  and the structure is symmetrical about  $VD$ .



- (i) Show that  $VD = 3 \cos \theta + 2 \sin \theta + 1.5$ .

[1]



- (ii) Express  $VD$  in the form  $R \cos(\theta - \alpha) + 1.5$ , where  $R > 0$  and  $\alpha$  is acute. [3]

$$\begin{aligned}
 \text{Let } 3 \cos \theta + 2 \sin \theta &= R \cos(\theta - \alpha) \\
 &= R \cos \theta \cos \alpha + R \sin \theta \sin \alpha \\
 R \cos \alpha &= 3 \quad \dots (1) \\
 R \sin \alpha &= 2 \quad \dots (2) \\
 \frac{(2)}{(1)}: \tan \alpha &= \frac{2}{3} \Rightarrow \alpha = \tan^{-1} \frac{2}{3} \approx 33.69^\circ \\
 (1)^2 + (2)^2: R^2 &= 3^2 + 2^2 \Rightarrow R = \sqrt{13} \\
 VD &= \sqrt{13} \cos(\theta - 33.69^\circ) + 1.5
 \end{aligned}$$

**Note:** Some students gave angle in radian mode instead.

- (iii) State the largest value of  $VD$  and find the corresponding value of  $\theta$ . [2]

$$\begin{aligned}
 \text{Large value of } VD &= \sqrt{13} + 1.5 \text{ or } 5.11 \\
 \text{when } \theta &= 33.69^\circ
 \end{aligned}$$

- (iv) Find the value(s) of  $\theta$  for which  $VD = 5$  m. [4]

$$\begin{aligned}
 \sqrt{13} \cos(\theta - 33.69^\circ) &= 3.5 \\
 \cos(\theta - 33.69^\circ) &= \frac{3.5}{\sqrt{13}} \\
 \text{Basic angle} &= \cos^{-1} \frac{3.5}{\sqrt{13}} \\
 &\approx 13.8978^\circ \\
 \theta - 33.69^\circ &= 13.8978^\circ, \quad -13.8978^\circ \\
 \theta &\approx 47.6^\circ, \quad 19.8^\circ
 \end{aligned}$$

**Note:** Many students missed out the answer,  $19.8^\circ$ .

4 (a) Solve the equation  $8e^{x+1} = \frac{15}{e^{x-1}} - 14e$ .

[4]

$$\begin{aligned}
 8e^{x+1} &= \frac{15}{e^{x-1}} - 14e \\
 8e(e^x) &= \frac{15}{e(e^x)} - 14e \\
 \text{Let } y &= e^x. \\
 8ey &= \frac{15}{ey} - 14e \\
 8y^2 + 14y - 15 &= 0 \\
 (4y-3)(2y+5) &= 0 \\
 y = \frac{3}{4} \quad \text{or} \quad y = -\frac{5}{2} \quad (\text{rej.}) \\
 e^x &= \frac{3}{4} \\
 x &= \ln \frac{3}{4} \approx -0.288
 \end{aligned}$$

(b) Solve the equation  $3\sin 2\theta = \cos \theta$  for  $-\pi \leq \theta \leq \pi$ .

[4]

$$\begin{aligned}
 3\sin 2\theta &= \cos \theta \\
 6\sin \theta \cos \theta &= \cos \theta \\
 6\sin \theta \cos \theta - \cos \theta &= 0 \\
 \cos \theta (6\sin \theta - 1) &= 0 \\
 6\sin \theta - 1 &= 0 \quad \text{or} \quad \cos \theta = 0 \\
 \sin \theta &= \frac{1}{6} \quad \theta = -\frac{\pi}{2}, \frac{\pi}{2} \\
 \theta &\approx 0.167, 2.79
 \end{aligned}$$

**Note:** Many students cancelled out  $\cos \theta$  and thus missed out one equation with 2 answers.

- 5 (a) Solve the equation  $\log_5 x + 1 = 2\log_x 5$ .

[4]

$$\log_5 x + 1 = 2\log_x 5$$

$$= 2\left(\frac{1}{\log_5 x}\right)$$

$$\text{Let } y = \log_5 x:$$

$$y + 1 = \frac{2}{y}$$

$$y^2 + y - 2 = 0$$

$$(y-1)(y+2) = 0$$

$$y = 1 \quad \text{or} \quad y = -2$$

$$\log_5 x = 1 \quad \text{or} \quad \log_5 x = -2$$

$$x = 5 \quad \text{or} \quad x = \frac{1}{25}$$

- (b) Solve the equation  $\log_9(2+x) = \log_{16} 4 - \log_{\frac{1}{3}} \sqrt{1-2x}$ .

[5]

$$\log_9(2+x) = \log_{16} 4 - \log_{\frac{1}{3}} \sqrt{1-2x}$$

$$\frac{\log_3(2+x)}{\log_3 9} = \frac{\log_2 4}{\log_2 16} - \frac{\log_3 \sqrt{1-2x}}{\log_3 \frac{1}{3}}$$

$$\frac{\log_3(2+x)}{2} = \frac{2}{4} - \frac{\frac{1}{2} \log_3(1-2x)}{-1}$$

$$\frac{\log_3(2+x)}{2} = \frac{1}{2} + \frac{1}{2} \log_3(1-2x)$$

$$\log_3(2+x) = 1 + \log_3(1-2x)$$

$$\log_3 \frac{2+x}{1-2x} = 1$$

$$\frac{2+x}{1-2x} = 3$$

$$2+x = 3-6x$$

$$7x = 1$$

$$x = \frac{1}{7}$$

- 6 (i) Show that  $\frac{d}{dx}[x\sqrt{3x-1}] = \frac{Ax+B}{\sqrt{3x-1}}$ , where  $A$  and  $B$  are constants. [4]

$$\begin{aligned}
 \frac{d}{dx}[x\sqrt{3x-1}] &= \sqrt{3x-1} + x \cdot \frac{1}{2}(3x-1)^{-\frac{1}{2}} \cdot 3 \\
 &= \sqrt{3x-1} + \frac{3}{2}x(3x-1)^{-\frac{1}{2}} \\
 &= (3x-1)^{-\frac{1}{2}} \left[ (3x-1) + \frac{3}{2}x \right] \\
 &= (3x-1)^{-\frac{1}{2}} \left[ \frac{9}{2}x - 1 \right] \\
 &= \frac{\frac{9}{2}x - 1}{\sqrt{3x-1}} \quad \text{or} \quad \frac{9x-2}{2\sqrt{3x-1}}
 \end{aligned}$$

- (ii) Hence or otherwise, find  $\int_1^2 \frac{9x-2}{\sqrt{3x-1}} dx$ , giving your answer correct to 3 significant figures. [4]

$$\begin{aligned}
 \int \frac{9x-2}{\sqrt{3x-1}} dx &= 2 \int \frac{9x-2}{2\sqrt{3x-1}} dx \\
 &= 2x\sqrt{3x-1} + c \\
 \int_1^2 \frac{9x-2}{2\sqrt{3x-1}} dx &= \left[ 2x\sqrt{3x-1} \right]_1^2 \\
 &= 2(2)\sqrt{3(2)-1} - 2(1)\sqrt{3(1)-1} \\
 &= 4\sqrt{5} - 2\sqrt{2} \\
 &\approx 6.12
 \end{aligned}$$

- 7 The function  $f$  is defined for  $x \neq -\frac{1}{3}$  and is such that  $f''(x) = 4e^{2x} + \frac{9}{(3x+1)^2}$ .

Given that  $f'(0) = -1$  and  $f(0) = 2$ , find an expression for  $f(x)$ .

[7]

$$\begin{aligned} f'(x) &= \int 4e^{2x} + \frac{9}{(3x+1)^2} dx \\ &= 2e^{2x} + \frac{9(3x+1)^{-1}}{(-1)(3)} + c \\ &= 2e^{2x} - \frac{3}{3x+1} + c \end{aligned}$$

$$\begin{aligned} f'(0) &= -1 \\ 2 - 3 + c &= -1 \\ c &= 0 \end{aligned}$$

$$f'(x) = 2e^{2x} - \frac{3}{3x+1}$$

$$\begin{aligned} f(x) &= \int 2e^{2x} - \frac{3}{3x+1} dx \\ &= e^{2x} - \ln(3x+1) + d \end{aligned}$$

$$\begin{aligned} f(0) &= 2 \\ 1 - \ln(1) + d &= 2 \\ d &= 1 \end{aligned}$$

$$f(x) = e^{2x} - \ln(3x+1) + 1$$



- 8 (a) The equation of a curve is  $y = \frac{1}{\sqrt{10x-1}} + \frac{1}{3}x$ , for  $x \neq \frac{1}{10}$ . Find the equation of the normal to the curve at the point  $\left(1, \frac{2}{3}\right)$ . [4]

$$y = \frac{1}{\sqrt{10x-1}} + \frac{1}{3}x$$

$$= (10x-1)^{-\frac{1}{2}} + \frac{1}{3}x$$

$$\frac{dy}{dx} = -\frac{1}{2}(10x-1)^{-\frac{3}{2}}(10) + \frac{1}{3}$$

$$= -\frac{5}{(10x-1)^{\frac{3}{2}}} + \frac{1}{3}$$

$$\left. \frac{dy}{dx} \right|_{x=1} = -\frac{5}{(10-1)^{\frac{3}{2}}} + \frac{1}{3}$$

$$= \frac{4}{27}$$


Gradient of normal at  $P = -\frac{27}{4}$

Eqn of normal at  $P$ :

$$y - \frac{2}{3} = -\frac{27}{4}(x-1)$$

$$y = -\frac{27}{4}x + \frac{89}{12} \quad \text{or} \quad 12y = 81x + 89$$

- (b) The diagram shows a cylindrical water tank with base radius 16 cm.



Water is flowing into the tank at a constant rate of  $4.8\pi k \text{ cm}^3/\text{min}$ , where  $k$  is a constant. At time  $t$  minutes, the depth of the water in the tank is  $h$  metres. There is a hole at point  $P$  at the bottom of the tank and a cork is used to stop water leakage. When the cork is unplugged, water will leak from the hole at a rate of  $6.4\pi \text{ cm}^3/\text{min}$ . Find an expression for the rate of change of  $h$  in terms of  $k$  after the cork is unplugged.

[4]

$$\frac{dV_{\text{in}}}{dt} = 4.8\pi k, \quad \frac{dV_{\text{out}}}{dt} = 6.4\pi$$

$$\frac{dV}{dh} = 256\pi$$

$$\begin{aligned} \frac{dh}{dt} &= \frac{dh}{dV} \times \frac{dV}{dt} \\ &= \frac{1}{256\pi} \times (4.8\pi k - 6.4\pi) \end{aligned}$$

$$\frac{dh}{dt} = \frac{3k}{160} - \frac{1}{40}$$

$$\text{or } \frac{dh}{dt} = \frac{3k - 4}{160}$$

- 9 (i) The expansion of  $(1+px)^n$ , where  $n > 0$ , by the binomial theorem is  $1+36x+66p^2x^2+kx^3+\dots$ . Find the value of  $n$ , of  $p$  and of  $k$ . [5]

$$\begin{aligned}(1+px)^n &= 1 + \binom{n}{1}(px) + \binom{n}{2}(px)^2 + \binom{n}{3}(px)^3 + \dots \\ &= 1 + np x + \binom{n}{2} p^2 x^2 + \binom{n}{3} p^3 x^3 + \dots\end{aligned}$$

Comparing  $x^2$ :

$$\begin{aligned}\binom{n}{2} &= 66 \\ \frac{n(n-1)}{2} &= 66 \\ n^2 - n - 132 &= 0 \\ (n-12)(n+11) &= 0 \\ n &= 12 \quad \text{or} \quad n = -11 \text{ (rej.)}\end{aligned}$$

Comparing  $x$ :

$$\begin{aligned}np &= 36 \\ 12p &= 36 \\ p &= 3\end{aligned}$$

Comparing  $x^3$ :

$$\begin{aligned}k &= \binom{n}{3} p^3 \\ &= \binom{12}{3} (3)^3 \\ &= 5940\end{aligned}$$

- (ii) Find the term independent of  $x$  in the expansion of  $(3+x)\left(2x-\frac{1}{x}\right)^8$ . [4]

$$\text{For } \left(2x - \frac{1}{x}\right)^8 :$$

$$\begin{aligned} T_{r+1} &= \binom{8}{r} (2x)^{8-r} \left(-\frac{1}{x}\right)^r \\ &= \binom{8}{r} 2^{8-r} (-1)^r x^{8-2r} \end{aligned}$$

$$\text{Let } 8 - 2r = 0 \Rightarrow r = 4$$

$$\begin{aligned} \text{Term independent of } x &= \binom{8}{4} 2^4 (-1)^4 \\ &= 1120 \end{aligned}$$

$$\text{For } (3+x)\left(2x-\frac{1}{x}\right)^8 :$$

$$\text{Term independent of } x = 3(1120)$$


$$= 3360$$

- 10 The table shows, to 2 decimal places, the population,  $P$ , of a certain kind of bacteria in a test-tube, measured at hourly intervals,  $t$ .

$t$	0.2	0.6	0.9	1.3	1.5
$P$	0.72	0.96	1.40	1.55	1.78

- (i) On the grid on the next page, plot  $\lg P$  against  $t$  and draw a straight line graph. [3]

$t$	0.2	0.6	0.9	1.3	1.5
$\lg P$	-0.14	-0.02	0.15	0.19	0.25



- (ii) Determine which value of  $P$ , in the table above, is the incorrect reading.  
Use the graph to estimate a value of  $P$  to replace the incorrect recording of  $P$ . [2]

Incorrect reading of $P = 1.40$
<p>From graph, when <math>t = 0.9</math>,</p> $\lg P = 0.075$ $P = 10^{0.075} \approx 1.19$

- (iii) Find the gradient of your straight line and hence express  $P$  in the form  $\frac{A^t}{10^{2k}}$ , where  $A$  and  $k$  are constants. [4]

<p><math>(0, -0.2), (1.2, 0.16)</math></p> $\text{Gradient} = \frac{-0.2 - 0.16}{0 - 1.2} \approx 0.3000$ <p><math>(\lg P) - \text{intercept} = -0.2</math></p> $\lg P = 0.3000t - 0.2$ $P = 10^{0.3000t - 0.2}$ $= \frac{(10^{0.3})^t}{10^{2(0.1)}} \quad \text{or} \quad \frac{(2)^t}{10^{2(0.1)}}$
---

- (iv) On the same diagram, draw the line representing  $P^{10} = 10^{3-t}$  and hence estimate the value of  $t$  for which  $\left(\frac{A^t}{10^{2k}}\right)^{10} = 10^{3-t}$ . [2]

$P^{10} = 10^{3-t}$ $10 \lg P = 3 - t$ $\lg P = -\frac{1}{10}t + \frac{3}{10}$ <p>Draw the straight line: <math>\lg P = -\frac{1}{10}t + \frac{3}{10}</math></p> <p>From graph, <math>t \approx 1.24</math></p>
---

END OF PAPER